

Variability of UV-vis-IR solar irradiance from GOME and SCIAMACHY for use in GCMs

J. Pagaran, M. Weber, K. Bramstedt, J. Burrows
IUP, Bremen, Germany

N. Krivova, and S. Solanki
MPI Katlenburg-Lindau, Germany

L. Floyd
NRL, USA



background image from spacecenter.dk



Outline

(input)
SSI
11-yr cycle




Problem: No direct 11-year SSI measurements

GCM
(wavelength dependent)

How to increase sensitivity of climate
to 11-yr solar cycle variability?

- Solar data from GOME and SCIAMACHY
- Modeling observed solar variability
- Validate model with other solar data
- Radiation intervals for GCMs
- Summary and Recommendation

Radiative heating by O₂ and O₃ absorption



O₂ absorption bands
 125 – 205 nm Schumann-Runge (SRB/SRC)
 205 – 243 nm Herzberg contin (Hz)

O₃ absorption bands
 243 – 278 nm Hartley band (Ha)
 278 – 363 nm Huggins bands (H)
 407 – 683 nm Chappuis band (C)

Objective:

Estimate the best realistic TOA incoming radiation at

(1) solar min and (2) solar max conditions --- 11-year spectral dependence

Simple irradiance model

SOLAR DYNAMO



SURFACE FIELD



SUNSPOTS

FACULAE

IRRADIANCE

TSI
solar
constant

SSI: UV-vis-IR

CLIMATE

SCIA solar data
240-1700 nm
2003-2004

$I(\lambda, t)$

$$= a(\lambda) \cdot P_{\text{Mg II}}(t) + b(\lambda) \cdot P_{\text{PSI}}(t) + P_{\text{OK}}(t)$$

$\Delta I(\lambda, t)$

anomaly term

Mg II ctw from Viereck et al. (2004)

PSI from Balmaceda et al. (2007)

choose best measurements

- * when instrument is stable
- * during least degradation
- * during active sun periods
- * of longer timeseries

Polynomial term

SOLAR DYNAMO



SURFACE FIELD



SUNSPOTS

FACULAE

IRRADIANCE

TSI
solar
constant

SSI: UV-vis-IR

CLIMATE

$$I(\lambda, t) = a(\lambda) \cdot P_{\text{Mg II}}(t) + b(\lambda) \cdot P_{\text{PS}}(t) + \boxed{\text{Pol}(t)}$$

Polynomial term

(polynomial degree kept minimum)

- * slowly varying background

- * instrument degradation

Most desired objects: $a(\lambda)$ and $b(\lambda)$

SOLAR DYNAMO



SURFACE FIELD



SUNSPOTS

FACULAE



IRRADIANCE

TSI
solar
constant



SSI: UV-vis-IR

CLIMATE

linear regression

$$I(\lambda, t)$$

$$= a(\lambda) \cdot P_{\text{Mg II}}(t) + b(\lambda) \cdot P_{\text{PSI}}(t) + \text{Pol}(t)$$




linearly scale 27-day SSI variability

to 11-year decadal changes

Step-by-step illustration


(4) Fit low deg polynomials

310 -- 320 nm





Modeling: step-by-step illustration

UV parametrization @ 310- 320 nm




$$\text{Ratio} = \frac{I_{\text{data}}(\lambda, t)}{\text{Pol}(t)}, \frac{I_{\text{data}}(\lambda, t) - \text{Res}}{\text{Pol}(t)}$$

Derived regression parameters




after parametrizing all 143 10-nm intervals

Reconstruction of daily irradiances




Reconstruction of daily spectra (1978-2007)



Validation: spectral aspect

4th March 1994


distant past:
outside time domain
where parameters
are derived



time slice: from 26 May 2003 to 22 Jun 2003


Validation: temporal aspect

at visible: 515 nm



Validation: temporal aspect

fig from Fröhlich





Validation: temporal aspect

$a(\lambda)$ $b(\lambda)$
daily TSI values

$$\text{TSI}_j = \int_{240 \text{ nm}}^{1670 \text{ nm}} I(\lambda, t_j) d\lambda$$


$t_j \in \text{satellite era}$

TSI record	Mean value (W m ⁻²)	Standard deviation (W m ⁻²)	Correlation with TIM (2003-2006)	Slope per year (W m ⁻²)
TIM	1360.98	0.579	1.	- 0.182
PMOD	1365.76	0.573	0.9964	- 0.134
ACRIM	1366.09	0.582	0.9829	- 0.261
Lean (model)	1365.95	0.479	0.9634	- 0.069
this work	1232.62	0.523	0.9186	- 0.038


Table of values from Lean, NRL

11-yr solar cycle UV-vis-IR variability


Vis-IR: 400-1700 nm



UV contribution to 11-year ΔTSI



vis-IR contribution to 11-year ΔTSI



Radiation intervals for GCMs

Band name Wavelength interval	No. of intervals	No. of modeled or calculated intervals			
		SCIA	GOME	SUSIM	SIP
Lyman α 121.5 nm	1 point	0	0	1	1
Schumann-Runge 125 – 205 nm	4 bands	0	0	4	4
Herzberg continuum 206 – 243 nm	15	0	0	15	15
Hartley bands 243 – 278 nm	10	10	10	10	10
Huggins bands 278 – 363 nm	18	18	18	18	18
Chappuis band 407 – 683 nm	1	1	1	0	0
Total	49	29	29	48	48
Method		A	A	B	B

SIP (Solar2000)


from Tobiska, SpaceWx

Method

A 11-yr extrapolation

B direct ratio

Radiation intervals for GCMs





Summary and Recommendation

- * A simple irradiance model parametrizes GOME and SCIAMACHY observed solar variations
 - using Mg II (faculae brightening) and PSI (sunspot darkening)
 - * Using the simple model,
 - we estimate the 11-yr solar cycle variability radiation intervals that are relevant for GCMs
- after doing model validations:
- * reconstructed daily spectra with SUSIM, SIM & SIP (**within 5%**)
 - * reconstructed TSI and correlate with TIM
(slope comparable with Lean's model)
 - * 11-year contribution to Δ TSI with SUSIM & SATIRE
(large difference at 300-400 nm)

We recommend that GCMs use our estimates of 11-year variability to improve sensitivity of solar cycle influence on climate.

Acknowledgements



SCIAMACHY is a collaboration between Germany, the Netherlands, and Belgium.

- * **Gerald Harder** and **Juan Fontenla** of LASP, University of Colorado for solar data from SIM/SORCE and from solar atmosphere model.
- * **W. Kent Tobiska** of Space Environment Technologies for Solar Irradiance Platform PG v2.33.

This work is supported by

(1) **DFG-CAWSES (Germany) SOLOZON**


„Solar variability and ozone interaction“



(2) **ISSI, Bern, Switzerland**

„Solar data, interpretation, and modeling“





Institut für Umweltphysik



Universität Bremen

