# Lecture 6 CME Energetics

February 6, 2017

## early observations and models of CMEs

CMEs observed during solar eclipses, and by Mauna Loa and SMM, showing their three-part structure. (Hundhausen et al, 80-90s, Low, Gibson et al., Low 1996).







*Figure 1.* Photograph of the 1974 total eclipse showing the white-light corona with a sketch of the large-scale coronal magnetic lines of force taken from Hundhausen (1994). The lines of force in the

Large Angle and Spectrometric COronagraph (LASCO, 1995 –) observes CMEs from earth view, with the FOV in solar radii: C1: 1.1 – 3 , C2: 1.5 – 6, C3: 3.5 – 30



#### modern coronagraphs observing CMEs

STEREO (2006 – ) carries EUVI (0 – 1.7), COR1 (1.5– 4), COR2 ( – 15), HI (– 1AU) instruments in a twin spacecraft A and B.



## the three-part white light CME





Mass and velocity are measured in white-light images.

$$E = \frac{1}{2}Mv^{2} + \frac{1}{2}Mv_{esc}^{2}\left(1 - \frac{R_{s}}{R}\right) + \frac{M}{m}kT$$

CME mass is estimated from the white-light images, where enhanced brightness is due to Thomson scattering by electrons (Vourlidas et al. 2000, 2006), ranging from 10<sup>13-16</sup> grams.



## height, velocity, and acceleration of CMEs



## height, velocity and acceleration of CMEs

#### Jmap from the twin spacecraft from different views



gure 3. Jmaps of the 2012 July 12–14 CME for each STEREO spacecraft along the propagation angle as measured by the raytrace method. The CME is cleared

# height, velocity and acceleration of CMEs

#### Forward fitting a CME structure (Thernisien et al. 2006, 2009)





Figure 1. Model fitting of CME ejecta and shock front. Images at 17:54 UT on 2012 July 12 from STEREO A COR2 (left) and STEREO B COR2 (right) are shown, along without (top) and with (bottom) the raytrace mesh. The green mesh shows the GCS fitting to the CME ejecta, whereas the red mesh shows the spheroid fitting to the CME shock front. A supplemental movie showing the CME and the mesh is provided.

(An animation and a color version of this figure are available in the online journal.)

## CME kinematics and flare emissions



#### CME kinematics and flare emissions



## CME kinematics and flare emissions





Model: Forbes & Priest Observation: Poletto & Kopp 1986, Fletcher et al. 2001, 2004, Qiu et al. 2002–2012, Isobe et al. 2002, 2005, Asai et al. 2002, 2004, ...

$$\Phi = \int B_R dA_R$$
$$E = v_{in} B_{in} = v_R B_R$$





Figure 1. Time history of the neutral-line electric field during the two-ribbon flare of 29 July, 1973, as computed from the Kopp-Poletto magnetic field model (solid curve) and from the method of Forbes and Priest (dashed curve).









# model prediction



Martens & Kuin 1989

model prediction



is shown in certain intervals before  $\sim 102,000$  s because two O-type nulls exist

### not all eruptions are success



Occasionally, failed filament eruptions are observed (Ji et al. 2003)



Magnetic properties of primordial CMEs and their surrounding from magnetic field extrapolations, e.g., NLFF, PFSS, used to test instability models (Liu et al. 2008 ...)







FIG. 2.—Profile of horizontal magnetic field above the erupted regions. The field is averaged along the magnetic neutral line of the active region where the eruption took place. This is a log-log plot. The green, red, and black curves represent FE, KI, and TI, respectively. The green and red curves with cross, star, diamond, and triangle symbols represent events 1–4 and events 5–8, while

FIG. 3.—Horizontal magnetic field above the erupted regions as a function of height. The field strength and height are both in log units. The height ranges from 42 to 840 Mm. The strength is computed by averaging the whole area of the active region where the eruption took place. Since we are going to examine behaviors of a magnetic field at high altitude that is comparable with the size of active region, the field averaged over the whole active region is

## coronal dimming and CME properties



Coronal dimming observed in SXR and EUV images indicates mass "loss" with an erupting CME, estimated to be  $10^{14}$ g (Hudson et al. 1996, Sterling & Hudson 1997)

## Coronal dimming and CME properties



#### Coronal dimming in EUV images

# Coronal dimming and CME properties





Dimming tracks the CME evolution and flare reconnection. Significant dimming occurs at the onset of flare reconnection (Cheng et al. 2016)



## Coronal dimming and CME properties



As CME rises, the emission measure (EM) decreases, and the corona becomes dimmer.

$$EM = \int n^2 dl$$
$$= \int n^2 \frac{dl}{dT} dT$$

DEM



#### Coronal dimming and CME properties



#### CME energetics with dimming analysis



Aschwanden



Figure 16. Log-log histograms of various physical parameters measured from the 399 CME events based on AIA/SDO data. Power-law fits are applied on the rightside tails of the distributions.

#### CME energetics with EUV and WL obs.

THE ASTROPHYSICAL JOURNAL, 831:105 (34pp), 2016 November 1

ASCHWANDEN THE ASTROPHYSICAL JOURNAL, 831:105 (34pp), 2016 November 1

ASCHWANDEN



Figure 18. Comparison of of CME masses (top panels), CME velocities (middle panels), and CME kinetic energies (bottom panels) between AIA and LASCO data sets, in form of (log-log) size distributions (left panels) and scatterplots (right panels). The diagonal solid line indicates identity, along with the means (dashed line) and standard deviations (dotted lines) of the logarithmic ratios.

(Aschwanden 2016)

energies are shown in the right column. Equality (dotted diagonal line), the mean (dashed line) and standard deviation (dotted line) of the energy ratios are indicated



- CME energy from coronagraph WL observations and EUV dimming observations
- Timing of CME acceleration, flare emission, and magnetic reconnection