### Space Weather Highlights 11 August 1997 - 17 August 1997

Solar activity was very low to low. Isolated C-class subflares occurred on 11 and 14 August.

Solar wind data were received from the WIND spacecraft a few hours per day. Velocities ranged from 350 - 440 km/sec through 12 August increasing to 570 km/sec on 13 August. Velocities gradually declined to 350 km/sec by the end of the period. Densities generally ranged from 03-11 p/cc, but increased to a peak of 18 p/cc on 12 August. Bz varied from minus 07 to plus 07 nT. Solar sector orientation was not readily discernible.

There were no significant proton enhancements observed at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes was at moderate to high levels during 15 - 17 August.

The geomagnetic field was mostly quiet to unsettled. However, the field became disturbed on 13 - 14 August with active levels reported at all latitudes and isolated minor storm periods at higher latitudes. Another brief disturbance occurred during the latter half of 17 August with active levels reported at all latitudes and isolated minor storm periods observed at higher latitudes.

#### Space Weather Forecast 20 August 1997 -15 September 1997

Solar activity is expected to be very low.

No significant proton enhancements are expected at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes is expected to be normal to moderate.

The geomagnetic field is expected to be predominately quiet to unsettled.



## Daily Solar Data

	Radio	Sun	Sunspot	X-ray				Flares				
	Flux	spot	Area	Background	X-	ray Fl	ux		Op	tical		
Date	10.7 cm	No. (	(10 <sup>-6</sup> hemi.)		C	M	X	S	1	2	3	4
11 August	80	53	220	B2.5	1	0	0	1	0	0	0	0
12 August	81	57	140	A9.5	0	0	0	3	0	0	0	0
13 August	82	61	180	A6.6	0	0	0	2	0	0	0	0
14 August	80	49	140	A7.2	1	0	0	1	0	0	0	0
15 August	78	60	120	A4.7	0	0	0	0	0	0	0	0
16 August	78	33	100	A5.7	0	0	0	0	0	0	0	0
17 August	76	23	60	A6.5	0	0	0	0	0	0	0	0

Daily Particle Data

			Dany Lan	ucie Daia					
		Proton Fluence otons/cm <sup>2</sup> -day		Electron Fluence (electrons/cm <sup>2</sup> -day-sr)					
Date	>1MeV	>10MeV	>100MeV	>.6MeV	>2MeV	>4MeV			
11 August	2.7E + 5	1.7E+4	4.0E+3		6.4E + 6	6			
12 August	3.1E+5	1.7E+4	3.9E+3		7				
13 August	4.8E + 5	1.8E+4	3.8E + 3		6.5E+6				
14 August	5.3E+5	1.7E+4	4.1E+3		1.1E+	7			
15 August	1.1E+6	1.8E+4	4.1E+3		1.1E + 3	8			
16 August	4.0E + 5	1.8E+4	4.1E+3		5.7E+7				
17 August	6.1E + 5	1.8E+4	4.3E+3		7				


Daily Geomagnetic Data

			oung C	Johnshon Dan		
	N	Iiddle Latitude		High Latitude	]	Estimated
	F	redericksburg		College		Planetary
Date	A	K-indices	A	K-indices	A	K-indices
11 August	10	2-4-2-2-2-2	5	2-1-0-4-1-0-0-0	9	2-4-2-2-3-2-2
12 August	7	1-1-1-2-2-1-3-3	*	*-0-1-3-0-0-0-1	5	1-1-1-2-1-2-2-3
13 August	17	3-3-4-4-3-2-3-3	3	2-1-1-1-2-1-1-0	12	3-2-4-3-1-2-2-3
14 August	18	3-5-4-2-2-3-3	9	2-4-4-1-0-2-1-0	17	3-5-5-2-1-2-2-3
15 August	8	1-2-3-3-2-2-1-1	4	2-1-2-0-1-1-1	8	1-2-3-3-2-2-1
16 August	8	3-2-3-2-2-1-1	6	1-1-4-3-0-0-0-0	9	3-3-3-3-2-1-2-1
17 August	10	2-1-2-3-3-3-2-2	4	2-0-1-1-2-2-1-2	9	2-1-2-3-3-3-3-2

Alerts and Warnings Issued

Date and Time of Issue (UT	Type of Alert or Warning	Date and Time of Event (UT)
11 Aug 601	K = 4 observed	11 Aug 03-06
14 Aug 601	K = 5 observed	14 Aug 03-06
15 Aug 1359	>2 MeV Electron Event ≥ 1000pft	ı 15 Aug 1259
17 Aug 0007	>2 MeV Electron Event in Progress ≥ 10	000pfu 15 Aug 1259
18 Aug 0009	>2 MeV Electron Event in Progress ≥ 10	000pfu 15 Aug 1259





## Twenty-seven Day Outlook

	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
20 Aug	74	10	3	03	76	5	2
21	72	5	2	04	78	5	2
22	72	5	2	05	78	8	3
23	72	5	2	06	78	8	3
24	72	5	2	07	78	8	3
25	72	5	2	08	78	5	2
26	72	8	3	09	78	10	3
27	72	10	3	10	78	10	3
28	72	8	3	11	78	8	3
29	72	5	2	12	78	8	3
30	72	10	2	13	76	8	3
31	72	5	2	14	76	5	2
01 Sep	72	5	2	15	74	5	2
02	74	5	2				



Energetic Event

	Time (UT)	X-ray	Optical Inform	nation	Peak	Sweep Freq
Date	1/2	Integ	Imp Location	Rgn	Radio Flux	Intensity
	Begin Max Max	Class Flux	Brtns Lat CMD	#	245 2695	II IV

#### No Event Observed

				Flare List			
						ptical	
		Time		X-ray	Imp /	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	#
11 August	0138	0145	0159	B3.2			
	B0917	U0918	0923	C1.1	SF	N22E02	8072
12 August	0554	0557	0604	B3.6	SF	N28E80	8074
	0931	0936	0941	B2.3	SF	N25E16	8071
	1240	1241	1248		SF	S20W38	8070
	1513	1516	1531	B1.2			
	1612	1616	1619	B1.3			
	2146	2151	2154	B1.8			
13 August	B0511	0517	0525		SF	S20W48	8070
C	0642	0647	0706	B2.1			
	1838	1843	1850	B3.3	SF	N26W03	8071
14 August	0129	0132	0147	C1.0	SF	N26W04	8071
15 August	0827	0834	0843	B1.4			
C	1605	1611	1619	B1.1			
	1903	1906	1909	B1.2			
16 August	0821	0826	0830	B1.4			
C	1343	1348	1358	B1.2			
17 August	0216	0224	0231	B1.8			
	1829	1834	1838	B3.9			



## Region Summary

	Location	1		Sunspot 6	Characteri	stics			Flares						
		Helio	Area	Extent	Spot	Spot	Mag		X-ray	/		0	ptica	1	_
Date (	° Lat ° CMD)		(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
	Re	gion 80	67												
_	N06E65	051	0010	00	AXX		Α								
03 Aug	N05E52	050	0000	00	AXX	001	A								
04 Aug	N05E39	050													
05 Aug	N05E26	050													
06 Aug	N05E13	050													
07 Aug	N05E00	050													
08 Aug	N05W13	050													
09 Aug	N05W26	050													
10 Aug	N05W39	050													
11 Aug	N05W52	050													
12 Aug	N05W65	050													
_								0	0	0	0	0	0	0	0
Died or	n Disk.														
Absolu	te heliograp	hic long	gitude: 05	60											
	Re	gion 80	68												
03 Aug	S22E33	069	0030	04	BXO	006	В				3				
04 Aug	S22E18	071	0020	01	HSX	002	A								
05 Aug	S21E06	070	0020	05	CSO	002	В								
06 Aug	S21W09	072	0010	01	HSX	001	A								
07 Aug	S20W22	072	0010	01	HRX	002	A								
08 Aug	S19W37	072	0000	00	AXX	001	A								
_	S19W50	072													
_	S19W63	072													
_	S19W76	072													
_	S19W89	072													
	,							0	0	0	3	0	0	0	0
Crossec	d West Lim	b.													



Absolute heliographic longitude:

070

## Region Summary - continued.

1	Sunspot Characteristics				Flares								
Helio	Area	Extent	Spot	Spot	Mag		X-ray	y	_	O	ptica	ıl	
Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
)													
023	0000	00	AXX	001	A								
024	0010	03	BXO	003	В								
025	0030	06	DSO	009	В								
024	0060	08	DAO	012	В								
023	0050	09	CSO	012	В								
024	0030	09	CSO	007	В								
025	0010	07	BXO	005	В								
024	0010	06	BXO	007	В				1				
024	0050	08	CAO	010	В				1				
023	0040	07	DSO	004	В								
024	0010	00	AXX	001	A								
024													
						0	0	0	2	0	0	0	0
b.													
hic lon	gitude: 02	23											
gion 80	071												
333	0060	04	CRO	004	В								
333	0010	06	BXO	003	В								
332	0010	04	BXO	004	В								
335	0030	06	CSO	006	В								
335	0030	05	CAO	008	В				1				
336	0030	05	CAO	006	В				1				
337	0010	01	HRX	002	Α	1			1				
338	0010	03	BXO	002	В								
338													
338													
						1	0	0	3	0	0	0	0
	Helio Lon   0	Helio   Area   Lon   (10 <sup>-6</sup> hemi)   O	Helio   Area   Extent   Lon   (10 <sup>-6</sup>   hemi)   (helio)   (helio	Helio	Helio	Helio	Helio	Helio	Helio	Helio	Helio	Helio	Helio

Still on Disk.

Absolute heliographic longitude: 335



Location	1	Sunspot Characteristics					Flares							
Location	Helio	Area	Extent	Spot	Spot	Mag		X-ray	У	11		ptica	1	
Date (° Lat ° CMD)	Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	С	M		S	1	-	3	4
Region 807.														
09 Aug N21E21	001	0000	00	AXX	001	A								
10 Aug N21E08	001													
11 Aug N21W05	001						1			1				
12 Aug N21W18	001													
13 Aug N21W31	001													
14 Aug N21W44	001													
15 Aug N21W57	001													
16 Aug N21W70	001													
							1	0	0	1	0	0	0	0
Died on Disk.														
Absolute heliograp	hic longi	tude: 00	)1											
Re	gion 807	'3												
11 Aug N14E74	281	0090	03	HSX	001	A								
12 Aug N15E61	281	0070	02	HSX	001	A								
13 Aug N14E49	280	0060	02	HSX	001	A								
14 Aug N14E36	279	0060	02	HSX	001	A								
15 Aug N15E23	279	0070	02	HSX	001	A								
16 Aug N15E08	281	0070	02	HSX	001	A								
17 Aug N15W04	280	0060	02	HSX	001	A								
a.u. 5							0	0	0	0	0	0	0	0
Still on Disk.		. 1 20												
Absolute heliograp	_	tude: 28	30											
Region 807-		0000	02	HCV	001									
11 Aug N26E75	280	0090	02	HSX	001	A				1				
12 Aug N26E63	279	0030	02	HSX	001	A				1				
13 Aug N26E52	277	0040	04	CSO	004	В								
14 Aug N27E38	277	0030	02	HSX	002	A								
15 Aug N27E27	275	0030	05	CSO	005	В								
16 Aug N27E11	278	0020	01	HRX		A								
17 Aug N27W02	278	0000	00	AXX	002	A	Λ	Ω	Ω	1	Ω	Ω	Λ	0
Still on Disk.							U	0	U	1	U	U	U	U
Absolute heliograp	hic longi	tude: 27	10											
0 1	gion 807		U											
15 Aug N29E46	256	0000	00	AXX	001	A								
16 Aug N29E31	258	0010	00		001	A								
17 Aug N29E18	258	0010	50	. 1. 1. 1	501	. 1								
171105 1127110	250						0	0	0	0	0	0	0	0
Still on Disk.							J	3	9	3	3	3	9	0
Absolute heliograp	hic longi	tude: 25	8											
and and and			-											




# Recent Solar Indices (preliminary) of the observed monthly mean values

of the observed monthly mean values												
			ot Numbers				o Flux	Geomagne				
-	Observed		Ratio	Smooth		**Penticton	Smooth	<u>Planetary</u>				
Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value			
					1995							
August	25.1	14.3	0.57	25.4	15.4	73.8	76.0	09	12.1			
September	16.5	11.8	0.72	22.0	13.4	72.0	74.8	13	11.8			
r			****			, =	,					
October	31.6	21.1	0.67	19.7	12.1	77.9	73.8	16	11.4			
November	15.7	09.0	0.57	18.5	11.4	74.2	73.2	09	10.7			
December	16.2	10.0	0.62	17.6	10.8	72.6	72.8	09	10.7			
December	10.2	10.0	0.02	17.0	10.0	72.0	12.0	0)	10.0			
					1996							
January	17.6	11.5	0.55	16.8	10.4	74.5	72.4	09	09.8			
February	09.1	04.4	0.48	16.2	10.1	71.5	72.2	10	09.8			
March	12.1	09.2	0.76	15.4	09.7	72.7	72.1	11	09.9			
17101011	12.1	07.2	0.70	10	07.7	, 2.,	, 2.1		0,,,			
April	08.5	04.8	0.60	13.6	08.6	69.3	71.6	11	09.7			
May	11.8	05.5	0.47	12.9	08.1*	72.1	71.4	07	09.5			
June	18.8	11.8	0.63	13.5	08.6*	69.6	71.4	05	09.4			
June	10.0	11.0	0.03	13.3	00.0	09.0	/1.0	03	07.4			
July	13.2	08.2	0.67	13.4	08.5*	71.2	72.0	07	09.3			
August	20.5	14.4	0.68	13.1	08.4*	72.4	72.1	09	09.4			
September	02.9	01.6	0.62	13.3	08.5*	69.4	72.3	15	09.3*			
•	02.2	01.0	0.79	140	00.0*	60.2	70.6	12	00.1*			
October	02.3	01.8	0.78	14.0	09.0*	69.2	72.6	13	09.1*			
November	26.7	18.6*	0.72*	15.4	10.0*	78.7	73.0	08	09.1*			
December	21.1	12.7*	0.60*	16.2	10.6*	77.8	73.3	07	09.3*			
					1997							
January	09.0	06.5*	0.72*	16.5	10.7*	74.0	73.5*	09	09.4*			
February	11.3	07.6*	0.67*			73.8		11				
March	14.4	08.8*	0.61*			73.5*		08*				
1viciren	1 1.1	00.0	0.01			73.3		00				
April	24.5	15.8*	0.64*			74.5*		10*				
May	28.6	18.5*	0.64*			74.6*		08*				
June	22.1	13.1*	0.59*			72.4*		08*				
3 (411)	<i></i> .1	13.1	0.07			, 4. 1		00				
July	17.0	10.5*	0.62*			71.2*		07*				

<sup>\*</sup>Preliminary estimates.

The lowest smoothed sunspot indices number for Cycle 21, RI = 12.3, occurred September 1986. The highest smoothed sunspot number for Cycle 22, RI=158.5, occurred July 1989. \*\* From June 1991 onward, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary

Week Beginning 11 Aug 1997


*Protons* plot contains the five minute averaged integral proton flux (protons/ cm<sup>2</sup>-sec-sr) as measured by GOES-9 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


*Electrons* plot contains the five minute averaged integral electron flux (electrons/ cm<sup>2</sup> -sec-sr) with energies greater than 2 MeV at GOES-9.

*Hp* plot contains the five minute averaged magnetic field H component in nanoteslas (nT) as measured by GOES-9. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis. *Kp* plot contains the estimated planetary 3-hour K-index (derived by the USAF 55<sup>th</sup> Space Weather Squadron) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA. These data are made available through cooperation from the Geological Survey of Canada (GSC) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. Hparallel is subject to a more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.








#### Weekly GOES Satellite X-ray and Proton Plots

Proton plot contains the five minute averaged integral proton flux (protons/cm<sup>2</sup>-sec-sr) as measured by GOES-9 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm<sup>2</sup>-sec-sr) at greater than 10 MeV.

X-ray plot contains five minute averaged x-ray flux (watts/m<sup>2</sup>) as measured by GOES 8 and 9 in two wavelength bands, .05 -.4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.



# Carrington Rotation 1925



*H-alpha Synoptic Chart*- - a partial and preliminary map of the sun in absolute heliographic coordinates (solar latitude and longitude). The Carrington Rotation serial number appearing at the top of the chart is a continuation of the sequence begun by R. Carrington on 09 November 1852. Dates along the top of the chart are the times of central meridian passage of the solar longitudes directly below those dates. Dates along the top of the chart are the times of central meridian passage of the solar longitudes directly below those dates. Features on the map are as follows: filaments (cross-hatched areas), filament channels (solid lines outside an active area), plage corridors (solid lines inside an active area), 01 and estimated neutral lines (dashed lines) are lines of magnetic polarity change (neutral lines), plages (dotted areas whose dot density is roughly equal to brightness), strong active regions (stippled areas overlaid with diagonal lines; source of x-ray flares class M2 or greater, or two or more class M1 flares), large sunspots (large dots), coronal holes (solid lines with tick marks directed toward center of coronal hole, from 1083 nm spectroheliograms), SWO region number showing the last 3 digits of a 4-digit number.

