Space Weather Highlights 18 August 1997 - 24 August 1997

Solar activity was very low. Isolated B-class X-ray flares occurred, a few of which were associated with small optical subflares.

Solar wind data were received from the WIND spacecraft a few hours per day. Velocities ranged from 330 - 450 km/sec. Particle densities generally ranged 03 - 13 p/cc with a peak to 18 p/cc noted on 24 August. Bz hovered about zero, ranging from minus 06 to plus 06 nT (GSM). Solar sector orientation was mostly away (phi angle near 135 degrees).

There were no significant proton enhancements observed at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes was at normal to moderate levels.

The geomagnetic field was mostly quiet to unsettled. However, a brief episode of active conditions occurred at all latitudes during 21/0300 - 0600UT.

Space Weather Forecast 27 August 1997 - 22 September 1997

Solar activity is expected to be very low to low.

No significant proton enhancements are expected at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes is expected to be normal to moderate.

The geomagnetic field is expected to be predominately quiet to unsettled.



Daily Solar Data

	Radio	Sun	Sunspot	X-ray				Flares				
	Flux	spot	Area	Background	X-	ray Fl	ux	Optical				
Date	10.7 cm	No.	(10 ⁻⁶ hemi.)		C	M	X	S	1	2	3	4
18 August	76	11	60	A2.7	0	0	0	0	0	0	0	0
19 August	74	11	60	A2.1	0	0	0	0	0	0	0	0
20 August	75	11	50	A1.6	0	0	0	0	0	0	0	0
21 August	75	11	60	A2.4	0	0	0	1	0	0	0	0
22 August	76	11	60	A5.5	0	0	0	0	0	0	0	0
23 August	77	11	20	A9.5	0	0	0	0	0	0	0	0
24 August	78	0	0	A6.6	0	0	0	1	0	0	0	0

Daily Particle Data

			Dany Lai	iicie Daia						
		Proton Fluence tons/cm ² -day			ectron Fluence rons/cm²-day					
Date	>1MeV	>10MeV	>100MeV	>.6MeV	>2MeV	>4MeV				
18 August	3.2E + 5	1.8E+4	4.0E+3		2.7E + 7	•				
19 August	2.6E + 5	1.9E+4	4.5E+3		•					
20 August	3.8E + 5	1.9E+4	4.5E+3		2.0E+7					
21 August	4.5E + 5	1.9E + 4	4.6E + 3		1.8E+7					
22 August	2.7E + 5	1.9E+4	4.7E+3		4.2E + 6	,)				
23 August	3.5E + 5	1.8E+4	4.5E+3		4.6E+6					
24 August	7.6E + 5	1.8E+4	4.4E+3		5.0E + 6	·				


Daily Geomagnetic Data

	N	Middle Latitude		High Latitude		Estimated
	Fredericksburg			College	-	Planetary
Date	A	K-indices	A	K-indices	A	K-indices
18 August	8	2-2-2-2-3-2-1	6	1-0-1-1-4-2-1-1	6	1-1-2-2-3-2-2
19 August	5	2-2-1-1-2-1-1	0	1-0-0-0-0-0-0	5	2-2-0-1-2-2-2-1
20 August	5	1-2-2-1-1-1-2-2	3	0-2-1-2-1-0-1-2	5	1-2-2-1-2-1-2-2
21 August	9	2-4-1-2-2-1-2-2	*	2-3-0-5-*-0-*-1	9	3-4-0-2-2-2-3
22 August	8	3-3-1-1-3-2-1-1	4	1-1-1-2-3-2-0-0	7	3-3-1-1-2-2-2
23 August	4	2-1-0-0-1-1-2-2	0	0-0-0-0-1-0-0-0	4	1-1-0-0-1-2-1-2
24 August	8	3-2-1-2-2-1-3-2	2	0-1-0-1-1-0-2-0	6	2-1-1-2-2-1-2-3

Alerts and Warnings Issued

Date and Time of Issu	ie (UT)	Type of Alert or Warning	Date and	Time of Event (UT)
18 Aug 0009	>2 N	MeV Electron Event in Progress ≥ 10	00pfu	17 Aug
21 Aug 0600		K = 4 observed	-	21 Aug 03-06





Twenty-seven Day Outlook

Radio Flux	Planetary	Largest		Radio Flux	Planetary L	argest	
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
27	86	12	3	10 Sep	78	15	3
28	86	10	3	11	78	8	3
29	86	10	3	12	78	10	3
30	86	10	3	13	76	10	3
31	86	8	3	14	76	5	2
01 Sep	84	5	2	15	74	5	2
02	84	5	2	16	74	5	2
03	82	5	2	17	74	10	3
04	82	5	2	18	76	8	3
05	82	8	3	19	76	5	2
06	80	8	3	20	78	5	2
07	78	8	3	21	80	5	2
08	78	5	2	22	82	8	3
09	78	15	3				



Energetic Event

	Time (U	T)	X-ray	Optical Inform	nation	Peak	Sweep Freq
Date		1/2	Integ	Imp Location	Rgn	Radio Flux	Intensity
	Begin Max	Max	Class Flux	Brtns Lat CMD	#	245 2695	II IV

No Event Observed

				Flare List			
					C	ptical	
		Time		X-ray	Imp /	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	#
18 August	1019	1023	1027	B1.1			
19 August	No Fla	res Obser	ved				
20 August	No Fla	res Obser	ved				
21 August	0433	0433	0438	B3.6	SF	N29W34	8074
22 August	No Fla	res Obser	ved				
23 August	1413	1434	1456	B3.1			
24 August	0626	0632	0642	B1.1			
3	1834	1835	1838	B2.4	SF	S23E72	
	2116	2151	2203	B3.5			

Region Summary

Location	ı		Sunspot	Characteri	stics					Fla	ares			
	Helio	Area	Extent	Spot	Spot	Mag	2	X-ray	7		О	ptica	1	
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	С	M	X	S	1	2	3	4
Re	gion 80	07 <i>1</i>												
08 Aug N21E62	333	0060	04	CRO	004	В								
09 Aug N26E49	333	0010	06	BXO	003	В								
10 Aug N26E37	332	0010	04	BXO	004	В								
11 Aug N25E20	335	0030	06	CSO	006	В								
12 Aug N26E07	335	0030	05	CAO	800	В				1				
13 Aug N26W07	336	0030	05	CAO	006	В				1				
14 Aug N26W22	337	0010	01	HRX	002	Α	1			1				
15 Aug N26W36	338	0010	03	BXO	002	В								
16 Aug N26W49	338													
17 Aug N26W62	338													
18 Aug N26W75	338													
19 Aug N26W88	338													
							1	0	0	3	0	0	0	0

Crossed West Limb.

Absolute heliographic longitude: 335



Region Summary - continued.

Location	1		Sunspot (Characteri	stics			Flares X-ray Optical C M X S 1 2 3						
	Helio	Area	Extent	Spot	Spot	Mag		X-ray	ay O I X S 1		ptica	ıl	_	
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
Re	gion 80	72												
09 Aug N21E21	001	0000	00	AXX	001	A								
10 Aug N21E08	001													
11 Aug N21W05	001						1			1				
12 Aug N21W18	001													
13 Aug N21W31	001													
14 Aug N21W44	001													
15 Aug N21W57	001													
16 Aug N21W70	001													
							1	0	0	1	0	0	0	0
Died on Disk.														
Absolute heliograp	hic long	gitude: 00	1											
Region 8073														
11 Aug N14E74	281	0090	03	HSX	001	A								
12 Aug N15E61	281	0070	02	HSX	001	A								
13 Aug N14E49	280	0060	02	HSX	001	A								
14 Aug N14E36	279	0060	02	HSX	001	A								
15 Aug N15E23	279	0070	02	HSX	001	A								
16 Aug N15E08	281	0070	02	HSX	001	A								
17 Aug N15W04	280	0060	02	HSX	001	A								
18 Aug N14W17	280	0060	02	HSX	001	A								
19 Aug N14W31	280	0060	02	HSX	001	A								
20 Aug N14W44	280	0050	02	HSX	001	A								
21 Aug N13W60	283	0060	01	HSX	001	A								
22 Aug N13W71	281	0060	02	HSX	001	A								
23 Aug N13W85	282	0020	01	AXX	001	A								
24 Aug N13W98	282													
C							0	0	0	0	0	0	0	0
Ctill on Dials														

Still on Disk.

Absolute heliographic longitude: 280



Region Summary - continued.

	Location	1		Sunspot (Characteri	stics		Flares							
		Helio	Area	Extent	Spot	Spot	Mag		X-ray	У		О	ptica	ıl	
Date	(°Lat°CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
	Region 807	4													
11 Au	g N26E75	280	0090	02	HSX	001	A								
12 Au	g N26E63	279	0030	02	HSX	001	A				1				
13 Au	g N26E52	277	0040	04	CSO	004	В								
14 Au	g N27E38	277	0030	02	HSX	002	A								
15 Au	g N27E27	275	0030	05	CSO	005	В								
16 Au	g N27E11	278	0020	01	HRX	001	A								
17 Au	g N27W02	278	0000	00	AXX	002	A								
18 Au	g N27W15	278													
19 Au	g N27W28	278													
20 Au	g N27W41	278													
21 Au	g N27W54	278									1				
22 Au	g N27W67	278													
23 Au	g N27W80	278													
24 Au	g N27W93	278													
								0	0	0	2	0	0	0	0
Still or	n Disk.														
Absol	ute heliograp	hic long	gitude: 27	' 8											
	Re	gion 80	75												
15 Au	g N29E46	256	0000	00	AXX	001	A								
16 Au	g N29E31	258	0010	00	AXX	001	A								
17 Au	g N29E18	258													
18 Au	g N29E05	258													
19 Au	g N29W08	258													
	g N29W21	258													
	g N29W34	258													
	g N29W47	258													
	g N29W60	258													
	g N29W73	258													
								0	0	0	0	0	0	0	0
Still or	n Disk.														



Absolute heliographic longitude: 258




Recent Solar Indices (preliminary) of the observed monthly mean values

		Sunsp	ot Numbers			mean values Radi	o Flux	Geomagnetic		
_	Observed	-	Ratio	Smooth	values	**Penticton	Smooth	_	Smooth	
Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value	
					1995					
August	25.1	14.3	0.57	25.4	15.4	73.8	76.0	09	12.1	
September	16.5	11.8	0.72	22.0	13.4	72.0	74.8	13	11.8	
October	31.6	21.1	0.67	19.7	12.1	77.9	73.8	16	11.4	
November	15.7	09.0	0.57	18.5	11.4	74.2	73.2	09	10.7	
December	16.2	10.0	0.62	17.6	10.8	72.6	72.8	09	10.0	
					1996					
January	17.6	11.5	0.55	16.8	10.4	74.5	72.4	09	09.8	
February	09.1	04.4	0.48	16.2	10.1	71.5	72.2	10	09.8	
March	12.1	09.2	0.76	15.4	09.7	72.7	72.1	11	09.9	
April	08.5	04.8	0.60	13.6	08.6	69.3	71.6	11	09.7	
May	11.8	05.5	0.47	12.9	08.1*	72.1	71.4	07	09.5	
June	18.8	11.8	0.63	13.5	08.6*	69.6	71.8	05	09.4	
July	13.2	08.2	0.67	13.4	08.5*	71.2	72.0	07	09.3	
August	20.5	14.4	0.68	13.1	08.4*	72.4	72.1	09	09.4	
September	02.9	01.6	0.62	13.3	08.5*	69.4	72.3	15	09.3*	
October	02.3	01.8	0.78	14.0	09.0*	69.2	72.6	13	09.1*	
November	26.7	18.6*	0.72*	15.4	10.0*	78.7	73.0	08	09.1*	
December	21.1	12.7*	0.60*	16.2	10.6*	77.8	73.3	07	09.3*	
					1997					
January	09.0	06.5*	0.72*	16.5	10.7*	74.0	73.5*	09	09.4*	
February	11.3	07.6*	0.67*			73.8		11		
March	14.4	08.8*	0.61*			73.5*		08*		
April	24.5	15.8*	0.64*			74.5*		10*		
May	28.6	18.5*	0.64*			74.6*		08*		
June	22.1	13.1*	0.59*			72.4*		08*		
July	17.0	10.5*	0.62*			71.2*		07*		

^{*}Preliminary estimates.

The lowest smoothed sunspot indices number for Cycle 21, RI = 12.3, occurred September 1986. The highest smoothed sunspot number for Cycle 22, RI=158.5, occurred July 1989. ** From June 1991 onward, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary

Week Beginning 18 Aug 1997


Protons plot contains the five minute averaged integral proton flux (protons/ cm²-sec-sr) as measured by GOES-9 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five minute averaged integral electron flux (electrons/ cm² -sec-sr) with energies greater than 2 MeV at GOES-9.

Hp plot contains the five minute averaged magnetic field H component in nanoteslas (nT) as measured by GOES-9. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis. *Kp* plot contains the estimated planetary 3-hour K-index (derived by the USAF 55th Space Weather Squadron) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA. These data are made available through cooperation from the Geological Survey of Canada (GSC) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. Hparallel is subject to a more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.








Weekly GOES Satellite X-ray and Proton Plots

Proton plot contains the five minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-9 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.


X-ray plot contains five minute averaged x-ray flux (watts/m²) as measured by GOES 8 and 9 in two wavelength bands, .05 -.4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.





Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona
These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.





Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona
These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.

