Space Weather Highlights 08 – 14 August 2005

SWO PRF 1563 16 August 2005

Solar activity was very low this period. Region 792 (N12, L=056, class/area, Dac/440 on 30 July) was the largest sunspot group on the disk, and rotated quietly around the west limb on 10 August. New Region 797 (S13, L=254, class/area, Dai/180 on 14 August) formed on the disk on 11 August, and showed steady growth during the period. The remainder of the disk and limb was quiet and stable.

Solar wind data were available from the NASA Advanced Composition Explorer (ACE) spacecraft during most of the summary period. Solar wind speed ranged from a high of near 600 km/s early on 08 August to a low of near 330 km/s early on 12 August. The period began with wind speeds elevated at near 600 km/s and the Bz component of the IMF weakly southward at –5 nT as the geomagnetic field was under the waning influence of a coronal hole high speed wind stream. Thereafter, wind speed steadily decreased to near 330 km/s by 12 August, while the IMF Bz generally did not vary much beyond +/- 5 nT through early on 13 August. At about 10/0600 UTC, transient flow in the solar wind increased wind speed slightly from about 420 km/s to about 465 km/s, while the IMF Bz transitioned from a northward orientation of about +3 nT to a southward orientation of between -5 to -9 nT for about 8 hours. Early on 13 August, a solar sector boundary crossing was observed in the ACE data. Solar wind speed gradually increased to 575 km/s by midday on 13 August, while the IMF Bz fluctuated between +17 and -12 nT during the same period. For the remainder of the summary period, the IMF Bz did not vary much beyond +/-5 nT, while solar wind speed ended the period at near 425 km/s.

No greater than 10 MeV proton events occurred during the summary period.

The greater than 2 MeV electron flux at geosynchronous orbit reached high levels on 08 and 09 August, and again on 12 August.

The geomagnetic field ranged from mostly quiet to unsettled levels, with one isolated active period observed at middle latitudes late on 13 August. At higher latitudes, an isolated active period was observed midday on 08 August, while a 12 hour period of minor to major storm levels were observed midday on 10 August.

Space Weather Outlook 17 August - 12 September 2005

Solar activity is expected be at very low to low levels.

No greater than 10 MeV proton events are expected.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at high levels on 17 - 22 August, 26 - 27 August, and 02 - 08 September.

The geomagnetic field is expected to range from quiet to major storm levels. Recurrent coronal hole high speed wind streams are expected to produce active to minor storm levels on 17 August, and active to major storm levels on 23 - 26 August. Otherwise, expect mostly quiet to unsettled conditions.



Daily Solar Data

				2000									
	Radio	Sun	Sunspot	X-ray	_			Flares					
	Flux	spot	Area	Area Background		-ray F	lux		Optical				
Date	10.7 cm	No.	(10 ⁻⁶ hemi.))	С	M	X	S	1	2	3	4	
08 August	86	56	370	B1.0	0	0	0	0	0	0	0	0	
09 August	83	51	360	A8.1	0	0	0	0	0	0	0	0	
10 August	76	34	190	A5.5	0	0	0	0	0	0	0	0	
11 August	76	35	220	A4.0	0	0	0	0	0	0	0	0	
12 August	76	47	210	A4.3	0	0	0	0	0	0	0	0	
13 August	75	33	110	A3.4	0	0	0	0	0	0	0	0	
14 August	75	34	190	A3.6	0	0	0	0	0	0	0	0	

Daily Particle Data

		oton Fluence ons/cm ² -day-si	r)	Electron Fluence (electrons/cm²-day-sr)
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV
08 August	6.4E+5	1.2E+4	2.7E+3	1.6E+9
09 August	5.0E + 5	1.2E+4	2.8E+3	1.4E+9
10 August	8.1E+5	1.3E+4	2.8E+3	4.9E+7
11 August	2.0E + 5	1.3E+4	3.3E+3	3.8E+7
12 August	2.5E+5	1.3E+4	3.2E+3	7.8E+7
13 August	3.2E + 5	1.3E+4	3.2E+3	5.4E+6
14 August	3.2E+5	1.3E+4	3.2E+3	9.0E+6

Daily Geomagnetic Data

				20011110		
	N	Middle Latitude		High Latitude		Estimated
	I	Fredericksburg		College		Planetary
Date	A K-indices		A	A K-indices		K-indices
08 August	6	2-2-2-1-2-2-1-2	9	2-2-2-1-4-2-1-2	8	2-2-3-1-2-2-1-3
09 August	8	2-3-2-2-0-3-2	9	2-2-3-3-3-3	10	3-3-2-2-2-3-3
10 August	9	2-1-3-3-3-2-1-1	39	3-2-6-6-5-1-0	22	3-2-4-5-5-3-2-2
11 August	2	1-0-1-1-1-0-0	4	1-0-2-2-3-1-0-0	6	1-1-2-1-2-2-2
12 August	3	1-1-1-0-1-0-1-2	6	1-1-2-3-3-0-1-1	6	1-1-1-2-2-2-3
13 August	14	3-3-2-3-1-3-4	11	2-3-2-3-3-1-2-3	16	3-4-2-3-3-2-3-3
14 August	6	2-2-1-1-1-1-2-1	9	3-2-2-3-3-1-1-1	10	4-4-2-2-2-2-2




Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
08 Aug 0515	ALERT: Electron 2MeV Integral Flux > 1000pfu	08 Aug 0500
09 Aug 0515	ALERT: Electron 2MeV Integral Flux > 1000pfu	<u>c</u>
10 Aug 0855	WARNING: Geomagnetic $K = 4$	10 Aug 0855 - 1500
10 Aug 0900	ALERT: Geomagnetic $K = 4$	10 Aug 0859
10 Aug 0912	WARNING: Geomagnetic $K = 5$	10 Aug 0915 - 1500
10 Aug 1035	ALERT: Geomagnetic $K = 5$	10 Aug 1035
10 Aug 1503	EXTENDED WARNING: Geomagnetic K=4	10 Aug 0855 - 2359
12 Aug 1126	ALERT: Electron 2MeV Integral Flux > 1000pfu	12 Aug 1105
13 Aug 0212	WARNING: Geomagnetic $K = 4$	13 Aug 0215 -1600
13 Aug 0412	ALERT: Geomagnetic $K = 4$	13 Aug 0411
13 Aug 0419	WARNING: Geomagnetic $K = 5$	13 Aug 0420 - 1600
14 Aug 0409	WARNING: Geomagnetic $K = 4$	14 Aug 0410 - 1600
14 Aug 0456	ALERT: Geomagnetic $K = 4$	14 Aug 0453



Twenty-seven Day Outlook



	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
17 Aug	75	20	5	31 Aug	100	12	3
18	75	15	4	01 Sep	100	10	3
19	80	8	3	02	95	12	3
20	80	7	2	03	90	12	3
21	80	7	2	04	85	8	3
22	80	7	2	05	85	10	3
23	85	15	4	06	80	10	3
24 Aug	90	25	6	07	75	8	3
25	95	20	5	08	75	8	3
26	100	15	3	09	75	15	4
27	105	8	33	10	75	12	3
28	105	12	3	11	75	10	3
29	105	12	3	12	75	12	3
30	100	10	3				



Energetic Events

			Bitter gett	ic Evenis			
	Time	X-ray	Opti	cal Information	l .	Peak	Sweep Freq
Date	1/2	Integ	Imp/	Location	Rgn	Radio Flux	Intensity
	Begin Max Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV

No Events Observed

				Flare List			
Date	Begin	Time Max	<u>End</u>	Optical X-ray Class.	Imp / Brtns	Location Lat CMD	Rgn
08 August	2019	2023	2025	B2.4			792
C	2310	2314	2317	B1.6			
09 August	0023	0028	0031	B2.7			792
G	1143	1148	1151	B1.7			796
	1204	1207	1209	B1.5			
	1841	1846	1853	B1.7			
10 August	0007	0010	0017	B1.4			
	0035	0039	0045	B2.1			792
	0240	0243	0247	B1.2			
	0312	0315	0319	B1.2			
11 August	1604	1610	1618	B1.3			795
12 August	1849	1857	1902	B1.2			797
13 August	1421	1440	1457	B1.7			
	1842	1845	1847	B1.6			797
14 August	1833	1839	1851	B1.1			797



Region Summary

Location	on			Character		<i>y</i>			F	lare	s			
Locatio	Helio	Area	Extent	Spot	Spot	Mag		X-ra		. —		Optic	al	
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi) (helio)	Class	Count	Class	C	M	X	S	1	2	3	4
Re	gion792	2												
28 Jul N12E78	057	0220	09	Dso	007	В	1	2		1				
29 Jul N11E66	055	0430	11	Ekc	028	Bg	1							
30 Jul N12E52	056	0440	09	Dac	019	Bgd	2		1	1	1	2		
31 Jul N12E38	057	0420	09	Dkc	040	Bgd		1						
01 Aug N12E25	057	0420	11	Ekc	034	Bgd	1	1		1	1			
02 Aug N12E12	057	0330	09	Dkc	043	Bgd	1			1				
03 Aug N11W01	057	0240	08	Dkc	026	Bg	2			1				
04 Aug N11W14	057	0180	07	Dki	018	Bg	1			1				
05 Aug N11W28	056	0160	07	Dso	018	Bg	1			1				
06 Aug N10W41	056	0130	05	Cso	009	Bg				1				
07 Aug N10W54	056	0110	06	Cso	800	В	1							
08 Aug N11W68	057	0120	05	Dso	005	В								
09 Aug N10W82	057	0110	04	Dao	003	В								
10 Aug N13W94	056	0030	01	Hax	001	A								
							11	4	1	8	2	2	0	0
Crossed West Lim														
Absolute heliograp	phic lon	gitude: 057												
Re	gion 79	4												
31 Jul S11E73	022	0060	03	Hax	002	A								
01 Aug S11E60	022	0110	06	Dao	007	В	1							
02 Aug S11E47	022	0150	06	Dao	008	В	1	1			1			
03 Aug S11E34	022	0210	08	Dsi	018	Bg	1	1		2	1			
04 Aug S11E19	022	0220	10	Dsi	029	Bg	1			2				
05 Aug S13E04	024	0230	10	Dai	024	Bg	1							
06 Aug S12W10	025	0220	08	Dho	011	Bg								
07 Aug S13W23	025	0170	08	Dso	010	Bg								
08 Aug S13W36	025	0120	05	Cso	005	В								
09 Aug S13W48	023	0140	03	Hax	003	A								
10 Aug S12W64	026	0130	04	Hax	002	A								
11 Aug S14W76	025	0140	02	Hax	002	A								
12 Aug S14W89	025	0120	02	Hax	001	A								_
							5	2	0	4	2	()	0	()

Crossed West Limb. Absolute heliographic longitude: 024



Region Summary - continued.

		K	<u> </u>			unuea						
Locatio		_				3.6		lares			1	_
(°I at °CMD)				-	.*	_						4
(Lat CIVID)	LOII	(10 Hellii) (Hello)	Class	Count	Class	C W X	<u>.s</u>	1		3	4
	Re	gion795										
N15E77	018	0060	02	Hax	002	Α						
ıg N15E64	018	0110	02	Hax	001	A						
ıg N15E51	018	0090	02	Hsx	003	Α						
ıg N15E38	018	0090	02	Hsx	002	Α						
ıg N13E27	014	0100	06	Cao	008	В						
ıg N13E14	014	0090	03	Hsx	002	A						
ıg N12E00	015	0070	04	Cso	004	В						
ıg N12W13	015	0060	04	Cso	004	В						
ıg N12W27	016	0050	02	Hsx	001	A						
ıg N11W40	015	0040	02	Hax	002	A						
ıg N15W55	017	0030	01	Hsx	001	A						
ıg N10W66	015	0040	01	Hax	001	A						
ıg N10W79	015	0030	03	Cro	004	В						
ıg N10W92	015	0030	01	Axx	001	A						
							$0 \ 0 \ 0$	0	0	0	0	0
ed West Lim	ıb.											
ute heliogra	phic lon	gitude: 015	i									
	-											
	_		03	Cac	002	D						
		0010	U3	Cso	002	D						
_												
•												
C		0070	0.4	D	005	D						
•												
•												
•		0070	05	Dao	003	В						
ıg S08W91	053						0 0 0	_	•	0	0	0
1777							0 0 0	0	0	O	0	U
	N15E77 1g N15E64 1g N15E51 1g N15E38 1g N15E38 1g N13E27 1g N13E14 1g N12E00 1g N12W13 1g N12W27 1g N11W40 1g N15W55 1g N10W66 1g N10W79 1g N10W92 1g S07E05 1g S07E05 1g S07W08 1g S07W21 1g S07W21 1g S07W34 1g S09W50 1g S08W64 1g S08W78 1g S08W91	N15E77 018 ag N15E64 018 ag N15E51 018 ag N15E38 018 ag N13E27 014 ag N13E14 014 ag N12E00 015 ag N12W13 015 ag N12W27 016 ag N12W27 016 ag N11W40 015 ag N15W55 017 ag N10W66 015 ag N10W79 015 ag N10W92 015 ag N10W92 015 ag N10W92 015 ag S07E05 051 ag S07E05 051 ag S07W21 051 ag S07W21 051 ag S07W34 051 ag S07W34 051 ag S09W50 052 ag S08W64 053 ag S08W78 053 ag S08W91 053	Cocation Helio Area (10-6 hemi helio Area (10-6 hemi helio Area (10-6 hemi helio hemi helio Area (10-6 hemi helio hemi helio	Location Helio Area Extent	Location Helio Area Extent Spot	Location Helio Area Extent Spot Spot	Cocation	Region 795	Location	Location	Location	Location

Crossed West Limb.

Absolute heliographic longitude: 051



Region Summary - continued.

Location	on		Sunspot	Characte					F	lares	S			
	Helio	Area	Extent	Extent Spot Spot I		Mag		X-ray				Optic	al	
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
Re	egion 797	7												
11 Aug S13E55	254	0040	02	Hax	002	Α								
12 Aug S13E42	254	0060	06	Dso	012	В								
13 Aug S13E29	254	080	07	Dai	012	В								
14 Aug S13E16	254	0180	08	Dai	013	В								
C							0	0	0	0	0	0	0	0
Still on Disk.														
Absolute heliogra	phic long	gitude: 254												
Re	egion 798	3												
14 Aug S05E54	216	0010	01	Hrx	001	A								
-							0	0	0	0	0	0	0	0
Still on Disk.														
Absolute heliogra	phic long	gitude: 216												
riosorate nenogra	pine long	511440. 210												




Recent Solar Indices (preliminary) of the observed monthly mean values

of the observed monthly mean values												
		Sunsp	ot Number	'S	-		Flux	Geomagne	etic			
	Observed	values	<u>Ratio</u>	Smooth	values	*Penticton	Smooth	Planetary	Smooth			
Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value			
				,	2003							
August	114.3	72.7	0.64	102.8	60.3	122.1	125.2	23	22.2			
September	82.6	48.8	0.59	100.7	59.8	112.3	123.7	18	21.8			
-												
October	118.9	65.5	0.55	96.6	58.4	153.1	121.8	35	21.1			
November	118.9	67.3	0.57	93.6	57.0	153.1	120.1	28	20.0			
December	75.4	46.5	0.62	91.4	55.0	115.1	118.0	16	18.6			
				,	2004							
January	62.3	37.7	0.61	87.9	52.0	114.1	116.3	22	18.1			
February	75.6	45.8	0.61	84.2	49.4	107.0	115.5	13	17.7			
March	81.0	49.1	0.61	80.9	47.2	112.2	114.6	14	16.9			
April	59.3	39.3	0.66	77.9	45.6	101.2	112.3	11	15.5			
May	77.3	41.5	0.54	74.1	43.9	99.8	109.2	8	14.3			
June	78.9	43.2	0.55	70.4	41.7	97.4	107.2	8	14.0			
T 1	07.0	51 0	0.50	60.2	40.0	110.5	1050	22	12.0			
July	87.8	51.0	0.58	68.3	40.2	118.5	105.9	23	13.8			
August	69.5	40.9	0.59	66.6	39.3	110.1	105.0	11	13.8			
September	50.0	27.7	0.55	63.7	37.6	103.1	103.7	10	13.6			
October	77.9	48.4	0.62	61.3	35.9	105.7	102.1	9	13.5			
November	70.5	43.7	0.62	60.0	35.4	113.2	101.5	26	14.1			
December	34.7	17.9	0.52	58.8	35.3	94.6	101.3	11	14.8			
					2005							
January	52.0	31.3	0.60	57.3	34.7	102.4	100.3	22	14.7			
February	45.4	29.1	0.64			97.3		11				
March	41.0	24.8	0.60			90.0		12				
April	41.5	24.4	0.59			85.9		12				
May	65.4	42.6	0.65			99.5		20				
June	59.8	39.6	0.66			93.7		13				
July	71.0	39.9	0.56			96.6		16				

NOTE: All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI = 120.8, occurred April 2000. *After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 08 August 2005

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W113) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm² –sec –sr) with energies greater than 2 MeV at GOES-12 (W75).


Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.







Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/ m^2) as measured by GOES 12 (W75) and GOES 10 (W135) in two wavelength bands, .05 - . 4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 (W113) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

