Space Weather Highlights 07 - 13 November 2005

SWO PRF 1576 15 November 2005

Solar activity was at very low to moderate levels. New Region 822 (S06, L=082, class/area, Cao/120 on 13 November) produced the majority of this period's activity. Late on 12 November and early on 13 November, this region produced two high level C-class flares. At 13/1451 UTC, Region 822 produced an impulsive M2.5 X-ray event; the first significant activity in almost two months. At the time of writing, Region 822 was rotating onto the disk where a detailed magnetic analysis was difficult. Other activity included a filament eruption near the central meridian late on 08 November. The remainder of the disk and limb was quiet and stable.

No greater than 10 MeV proton events were observed this period.

The greater than 2 MeV electron flux at geosynchronous orbit was at high levels on 07 - 11 November.

The geomagnetic field, at middle latitudes, ranged from quiet to unsettled levels with an isolated active period early on 13 November. At high latitudes, the field was generally quiet to unsettled through early on 12 November. Thereafter, the field was generally at active to major storm levels. Solar wind speed ranged from a high of near 700 km/s early on 07 November to a low of about 340 km/s late on 10 November. The period began with wind speed elevated at near 700 km/s, but decaying, and the Bz component of the IMF not varying much beyond +/- 3 nT. These conditions persisted through late on 10 November. Early on 11 November, a weak shock was detected in the ACE SWEPAM data as wind speed slowly elevated to near 400 km/s, and the IMF Bz fluctuated weakly to +/- 7 nT. This shock was believed to be associated with the filament eruption late on 08 November. The period ended with wind speed near 400 km/s and the IMF Bz not varying much beyond +/- 5 nT.

Space Weather Outlook 16 November - 12 December 2005

Solar activity is expected to be at low to moderate levels through 25 November when Region 822 is due to rotate around the west limb. Thereafter, very low to low activity is expected.

No greater than 10 MeV proton events are expected.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at high levels on 16 - 17 November, 01 - 08 December, and 12 December.

The geomagnetic field is expected to range from quiet to minor storm levels. Unsettled to active levels are possible on 21 November, while unsettled to minor storm levels are possible on 30 November – 01 December due to recurrent coronal hole high speed wind streams. Otherwise, expect quiet to unsettled conditions.



Daily Solar Data

				2	••••							
	Radio	Sun	Sunspot	X-ray	Fl			Flares				
	Flux	spot	Area	Background	X	-ray F		Optical				
Date	10.7 cm	No.	(10 ⁻⁶ hemi.))	С	M	X	S	1	2	3	4
07 November	r 79	31	180	A5.2	0	0	0	0	0	0	0	0
08 Novembe	r 79	38	120	A4.9	0	0	0	0	0	0	0	0
09 Novembe	r 78	13	30	A4.1	1	0	0	0	0	0	0	0
10 November	r 78	0	0	A3.1	0	0	0	0	0	0	0	0
11 Novembe	r 79	0	0	A2.1	0	0	0	0	0	0	0	0
12 Novembe	r 83	11	120	B1.2	2	0	0	0	0	0	0	0
13 November	r 88	16	120	B2.4	6	1	0	3	0	0	0	0

Daily Particle Data

		oton Fluence	Electron Fluence					
	(prote	ons/cm ² -day-sı	<u>;) </u>	(electrons/cm ² -day-sr)				
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV				
07 November	1.1E+6	1.5E+4	3.3E+3	1.8E+8				
08 November	1.1E+6	1.5E+4	3.3E+3	2.4E + 8				
09 November	1.4E+6	1.5E+4	3.6E + 3	3.8E+8				
10 November	1.4E+6	1.5E+4	3.5E+3	3.4E+8				
11 November	1.1E+6	1.6E+4	3.5E+3	1.1E+8				
12 November	8.1E + 5	1.6E+4	3.8E+3	2.6E+7				
13 November	3.8E + 5	1.6E+4	3.7E+3	1.2E+7				

Daily Geomagnetic Data


	Buily Geomagnette Build											
	N	Iiddle Latitude	I	High Latitude		<u>Estimated</u>						
	F	redericksburg		College		Planetary						
Date	Α	K-indices	A	K-indices	Α	K-indices						
07 November	6	3-1-3-2-1-1-0	7	3-1-3-2-2-1-2-0	6	3-1-2-1-1-1-1						
08 November	2	1-0-1-0-1-1-0-1	3	2-0-2-1-0-1-0-1	3	1-0-1-1-1-0-1						
09 November	1	2-0-0-0-0-1-0	1	1-0-1-1-0-0-0-0	3	2-0-0-0-0-1-0-1						
10 November	1	0-0-1-0-0-0-1	2	0-0-1-1-0-0-0-2	2	0-0-1-0-0-0-1-1						
11 November	3	1-2-0-1-1-1-1	4	0-1-0-3-2-2-1-0	5	1-2-1-1-2-2-1-1						
12 November	7	2-3-2-2-2-1-1	33	1-5-4-5-6-5-2-1	10	2-3-2-2-3-3-2-1						
13 November	12	3-4-3-2-2-1-3	37	1-2-3-6-6-6-4-2	14	3-4-3-3-3-2-3						

Alerts and Warnings Issued

	9	
Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
07 Nov 0848	ALERT: Electron 2MeV Integral Flux > 1000pfu	07 Nov 0820
08 Nov 0515	ALERT: Electron 2MeV Integral Flux > 1000pfu	08 Nov 0500
09 Nov 0515	ALERT: Electron 2MeV Integral Flux > 1000pfu	09 Nov 0500
10 Nov 0515	ALERT: Electron 2MeV Integral Flux > 1000pfu	10 Nov 0500
11 Nov 1002	ALERT: Electron 2MeV Integral Flux > 1000pfu	11 Nov 0940
13 Nov 0418	WARNING: Geomagnetic $K = 4$	13 Nov 0418 - 1500
13 Nov 0420	ALERT: Geomagnetic $K = 4$	13 Nov 0420



Twenty-seven Day Outlook



	Radio Flux	Planetary	Largest		Radio Flux	Radio Flux Planetary		
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Largest Kp Index	
16 Nov	100	10	3	30	75	20	5	
17	100	8	3	01 Dec	75	20	5	
18	100	8	3	02	75	12	4	
19	100	8	3	03	75	8	3	
20	100	8	3	04	75	8	3	
21	95	15	4	05	75	5	2	
22	95	10	3	06	75	8	3	
23 Nov	90	8	3	07	75	10	3	
24	85	8	3	08	75	10	3	
25	85	8	3	09	80	12	3	
26	90	8	3	10	85	10	3	
27	75	12	4	11	85	8	3	
28	75	8	3	12	85	8	3	
29	75	10	3					



Energetic Events

				2110180						
	Time		X-ray	Optical Information			Peak	Sweep Freq		
Date	Date		Integ	Imp/	Location	Rgn	Radio Flux	Intensity		
	Begin Max	Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV		
13 Nov	1429 1451	1456	M2.5 .016			822				

Flare List

	Flare List										
				Optical			_				
Data	Begin	Time Max	End	X-ray Class.	Imp / Brtns	Location Lat CMD	Rgn				
Date					Diuis	Lat CIVID					
07 November	0232	0235	0239	B1.0			820				
	0356	0400	0410	B1.0			820				
	0740	0805	0821	B2.2			820				
	1648	1655	1707	B1.6							
	1934	1938	1940	B1.6			819				
	2052	2059	2107	B4.3			820				
	2229	2234	2237	B3.4			819				
08 November	0225	0237	0254	B2.4			820				
09 November	0200	0205	0209	B2.5			820				
	0241	0301	0306	C1.2			820				
10 November	No Fla	resObserv	ved								
11 November	No Fla	resObserv	ved								
12 November	0311	0317	0342	B3.4							
	0414	0424	0439	B3.3			822				
	0736	0739	0741	B3.2			822				
	0941	1006	1018	B9.9			822				
	1332	1340	1349	B5.0			822				
	1418	1458	1603	C2.5			822				
	1817	1828	1837	B3.8			822				
	1843	1846	1850	B4.6			822				
	1917	1921	1923	B5.6			822				
	1925	1933	1937	C9.5			822				
	1958	2003	2007	B6.9			822				
13 November	0013	0036	0045	C9.5			822				
	0205	0225	0240	B7.6			822				
	0309	0406	0413	C1.9			822				
	0520	0522	0523	C6.7	Sf	S09E77	822				
	0602	0603	0607	C2.7	Sf	S09E82	822				
	0717	0717	0722		Sf	S09E82	822				
	1241	1249	1257	C2.7			822				
	1414	1417	1419	B3.8							
	1429	1451	1456	M2.5			822				
	1923	1926	1928	B3.9							
	2001	2009	2020	B7.4			822				
	2119	2158	2216	C2.8			822				



Region Summary

		y													
	Locatio	Helio	Area		Characte		Mag		V ro		lare		Intio	ol.	
Date	(°Lat°CMD)		(10 ⁻⁶ hemi)	Extent (helio)	Spot Class	Spot Count	Mag Class	\overline{C}	X-ra M	X	. <u>-</u>	1	Optic 2	<u>аг</u> 3	4
		gion 81													
31 O	ct S09E26	293	0020	03	Cso	003	В								
	ov S09E12	294	0030	04	Cso	006	В								
	ov S09W01	294	0030	06	Bxo	007	В								
	ov S09W14	294	0010	01	Axx	001	A				1				
	ov S07W28	294	0010	01	Axx	001	A								
	ov S07W41	294													
	ov S11W57	297	0000	01	Axx	001	A								
	ov S12W72	299	0120	05	Cao	003	В								
	ov S09W86	300	0030	04	Bxo	002	В								
								0	0	0	1	0	0	0	0
Cross	sed West Lim	ıb.													
Abso	olute heliogra	ohic lon	gitude: 294												
	$R \rho$	gion 82	20												
05 N	ov S16W27	280	0050	04	Dao	008	В								
	ov S17W41	281	0070	09	Dao	013	В	1							
	ov S17 W 11	281	0060	06	Cao	008	В	1							
	ov S17W69	283	0080	07	Cao	004	В								
	ov S16W81	281	0030	05	Bxo	003	В								
	ov S16W94	281	0020	0.0	Dirio	002	2								
1011	0,010,00	201						1	0	0	0	0	0	0	0
Cross	sed West Lim	ıb.													
Abso	olute heliograj	phic lon	gitude:280												
	Re	gion 82	21												
08 N	ov S14W19	233	0010	02	Axx	002	A								
	ov S13W32	233	0010	~ _		002									
	ov S13W45	233													
	ov S13W58	233													
	0,210,,00							0	0	0	0	0	0	0	0
Still	on Disk.														
	olute heliogra	ohic lon	gitude:233												
			· ·												
10 N		gion 82 080		02	Llow	001	٨	2							
	ov S04E81 ov S06E66	080	0120 0120	02 08	Hsx	001	A B	2 6	1		2				
13 IN	OV 200E00	082	0120	Uð	Cao	UUO	D	8	1	0	3	Λ	0	Λ	0
Q+i11 .	on Disk.							O	1	U	J	U	U	U	U
	oli Disk. olute heliograj	hic lon	oitude 27												
AUSC	nuic nenograj	one ion	Z1100C.02												




Recent Solar Indices (preliminary) of the observed monthly mean values

	of the observed monthly mean values											
			Sunsp	ot Number	·s	-	Radio	Flux	Geomagne	etic		
		Observed	values	Ratio	Smooth	values	*Penticton	Smooth	Planetary	Smooth		
_	Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value		
2003												
	November	118.9	67.3	0.57	93.6	57.0	153.1	120.1	28	20.0		
	December	75.4	46.5	0.62	91.4	55.0	115.1	118.0	16	18.6		
	December						110.1	110.0	10	10.0		
	т	<i>(</i> 2.2	27.7	0.61		2004	1141	1160	22	10.1		
	January	62.3	37.7	0.61	87.9	52.0	114.1	116.3	22	18.1		
	February	75.6	45.8	0.61	84.2	49.4	107.0	115.5	13	17.7		
	March	81.0	49.1	0.61	80.9	47.2	112.2	114.6	14	16.9		
	A	50.2	20.2	0.66	77.0	15.6	101.2	112.2	11	155		
	April	59.3	39.3	0.66	77.9	45.6	101.2	112.3	11	15.5		
	May	77.3	41.5	0.54	74.1	43.9	99.8	109.2	8	14.3		
	June	78.9	43.2	0.55	70.4	41.7	97.4	107.2	8	14.0		
	July	87.8	51.0	0.58	68.3	40.2	118.5	105.9	23	13.8		
	August	69.5	40.9	0.58	66.6	39.3	110.5	105.9	11	13.8		
	_						103.1					
	September	50.0	27.7	0.55	63.7	37.6	103.1	103.7	10	13.6		
	October	77.9	48.4	0.62	61.3	35.9	105.7	102.1	9	13.5		
	November		43.7	0.62	60.0	35.4	113.2	101.5	26	14.1		
	December	34.7	17.9	0.52	58.8	35.3	94.6	101.3	11	14.8		
					,	2005						
	January	52.0	31.3	0.60	57.3	34.7	102.4	100.3	22	14.7		
	February	45.4	29.1	0.64	56.4	34.0	97.3	98.5	11	14.6		
	March	41.0	24.8	0.60	55.8	33.6	90.0	97.2	12	15.3		
	March	11.0	21.0	0.00	33.0	33.0	70.0	71.2	12	13.3		
	April	41.5	24.4	0.59	52.6	31.7	85.9	95.5	12	15.7		
	May	65.4	42.6	0.65			99.5	, , ,	20			
	June	59.8	39.6	0.66			93.7		13			
	o dillo	27.0	27.0	0.00			,,,,		10			
	July	71.0	39.9	0.56			96.6		16			
	August	65.6	36.4	0.55			90.7		16			
	September		22.1	0.56			90.8		21			
	-											
	October	13.0	8.5	0.65			76.7		7			

NOTE: All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI= 120.8, occurred April 2000. *After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 07 November 2005

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W112) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm² –sec –sr) with energies greater than 2 MeV at GOES-12 (W76).

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/m²⁾ as measured by GOES 12 (W76) and GOES 10 (W135) in two wavelength bands, .05 - .4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 (W112) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

