Space Weather Highlights 31 July – 06 August 2006

SEC PRF 1614 08 August 2006

Solar activity was at very low levels. Through 03 August, the disk was only populated by small, magnetically weak spot groups. Since the 3rd, the disk was spotless.

No greater than 10 MeV proton events were observed.

The greater than 2 MeV electron flux at geosynchronous orbit was at normal levels.

The geomagnetic field was at quiet to unsettled levels at middle latitudes, while high latitudes observed quiet to minor storm conditions. Solar wind speed ranged from a low of near 380 km/s early on 31 July to a high of near 640 km/s midday on 01 August. The Bz component of the IMF did not vary much beyond +/- 3 nT for the majority of the period. Early on 31 July, wind speed and temperature began a slow increase, all in advance of a recurrent coronal hole wind stream. By midday on the 31st, the IMF Bz was fluctuating between +/-11 nT and remained so through the balance of the 31st. By midday on 01 August, wind speed peaked at about 640 km/s, while the IMF Bz relaxed some, not varying much beyond +/- 5 nT through the end of the 1st. As a result, at middle latitudes the geomagnetic field was at quiet to unsettled levels through midday on 01 August, while high latitudes experienced quiet to minor storm conditions through midday on 02 August. Thereafter and through the end of the summary period, the field was at mostly quiet levels.

Space Weather Outlook 09 August – 04 September 2006

Solar activity is expected to be at very low to low levels.

No greater than 10 MeV proton events are expected.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at high levels on 09-10 August.

The geomagnetic field is expected to be mostly quiet to unsettled for the majority of the forecast period. Active periods are possible on 24 and 28 August, while active to minor storm periods are possible on 03 September, all due to recurrent coronal hole high speed wind streams due to rotate into geoeffective positions.



Daily Solar Data

				2 1111 50								
	Radio	Sun	Sunspot	X-ray	_		Flares					
	Flux	spot	Area	Background	X	-ray F	lux		Or	otical		
Date	10.7 cm	No.	(10 ⁻⁶ hemi.))	С	M	X	S	1	2	3	4
31 July	72	25	30	A1.6	0	0	0	0	0	0	0	0
01 August	73	11	10	A1.4	0	0	0	0	0	0	0	0
02 August	72	22	50	A1.4	0	0	0	0	0	0	0	0
03 August	71	23	60	A2.4	0	0	0	0	0	0	0	0
04 August	70	0	0	A1.4	0	0	0	0	0	0	0	0
05 August	70	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
06 August	70	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0

Daily Particle Data

		oton Fluence ons/cm ² -day-si	r)	Electron Fluence (electrons/cm²-day-sr)
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV
31 July	2.3E+6	1.6E+4	3.9E+3	3.3E+6
01 August	6.9E + 5	1.6E+4	3.5E+3	2.6E+6
02 August	6.3E + 5	1.6E+4	3.5E+3	1.0E+7
03 August	4.6E + 5	1.6E+4	3.5E+3	2.3E+7
04 August	8.9E + 5	1.6E+4	3.6E+3	3.4E+7
05 August	8.9E + 5	1.7E+4	3.6E + 3	3.0E+7
06 August	1.3E+6	1.7E+4	3.6E+3	2.7E+7

Daily Geomagnetic Data


				oning neme Dana		
	M	Iiddle Latitude]	High Latitude		Estimated
	F	redericksburg		College]	Planetary
Date	A	K-indices	A	K-indices	A	K-indices
31 July	11	2-3-3-3-1-2-2	16	3-2-2-4-5-3-1-2	12	2-3-3-3-2-2-3
01 August	9	2-3-3-3-2-2-1-1	25	3-4-4-5-5-4-2-1	12	2-4-4-3-2-2-1-2
02 August	7	3-2-2-2-1-1-2	12	2-2-3-4-4-1-1-1	10	3-2-3-3-2-1-1-3
03 August	5	3-2-1-1-1-0-1-2	5	3-2-1-1-2-0-1-1	6	3-2-0-1-0-1-2-2
04 August	2	0-2-0-0-0-0-1-1	1	0-1-0-0-0-0-0-1	3	1-1-0-0-0-0-1-1
05 August	2	1-1-1-0-1-0-1-1	1	1-1-1-0-0-0-0-0	4	2-1-1-0-1-1-1
06 August	2	1-0-1-0-1-0-0-1	2	1-1-0-0-1-0-0-1	4	1-1-1-0-1-2-2-1

Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
01 Aug 0504	WARNING: Geomagnetic K=4	01 Aug 0505 - 1600
01 Aug 0506	ALERT: Geomagnetic K=4	01 Aug 0504
01 Aug 0854	ALERT: Geomagnetic K=4	01 Aug 0851



Twenty-seven Day Outlook



-	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
09 Aug	70	10	3	23 Aug	70	8	3
10	70	8	3	24	70	15	3
11	75	8	3	25	70	5	2
12	75	5	2	26	70	8	3
13	75	5	2	27	70	10	3
14	75	5	2	28	70	15	4
15	75	5	2	29	70	10	3
16	75	5	2	30	70	8	3
17	75	5	2	31	70	5	2
18	75	5	2	01 Sept	70	5	2
19	75	5	2	02	70	10	3
20	75	5	2	03	70	20	5
21	75	5	2	04	70	12	4
22	70	5	2				



Energetic Events

Time				X-ray	X-ray Optical Information				Sweep Freq
Date	Date ½		Integ	Imp/	Location	Rgn	Radio Flux	Intensity	
	Begin	Max	Max	Class Flux	Brtns	Lat CMD	#	245 2695	<u>II IV</u>
No Ever	nts Obser	ved							

				Flare List			
<u>Date</u>	Begin	Time Max	End	Optical X-ray Class.	Imp / Brtns	Location Lat CMD	Rgn
31 July	2010	2022	2029	B2.1			901
01 August	1031	1036	1040	B1.1			
02 August	No Fla	res Obser	ved				
03 August	No Fla	res Obser	ved				
04 August	0127	0134	0149	B1.8			
_	0326	0334	0353	B1.4			902
05 August	No Fla	res Obser	ved				
06 August	No Fla	res Obser	ved				

Region Summary

			8	<i>J</i>			
	Location	Sunspo	t Characteristics				
			Flares				
	Helio	Area Extent	Spot Spot	Mag	X-ray	Optical	
Date	(°Lat°CMD) Lon	(10 ⁻⁶ hemi) (helio)	Class Count	Class	C M X	S 1 2 3 4	

Region 901

22 Jul	N06E70	006	0040	06	Cso	006	В
23 Jul	N06E56	007	0130	08	Dao	011	В
24 Jul	N05E42	800	0110	08	Dao	009	В
25 Jul	N06E28	009	0100	08	Dao	006	В
26 Jul	N06E15	009	0100	09	Cao	010	В
27 Jul	N05E00	011	0070	09	Cso	013	В
28 Jul	N05W12	009	0070	09	Cso	007	В
29 Jul	N06W26	009	0080	09	Cso	009	В
30 Jul	N06W43	013	0010	01	Hsx	001	A
31 Jul	N06W54	011	0020	08	Dso	004	В
01 Aug	3 N06W67	011					
02 Aug	N08W73	003	0020	01	Axx	001	A
03 Aug	3 N08W86	003	0030	06	Bxo	002	В

0 0 0 0 0 0 0 0

Crossed West Limb.

Absolute heliographic longitude: 011



Region Summary-Continued

	Location	n			Character	ristics									
					Flares										
		Helio	Area	Extent	Spot	Spot	Mag	_	X-ra		. —)ptic	al	
Date (° La	at ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	С	M	X	S	1	2	3	4
	Re_{i}	gion 90.	2												
30 Jul S1	0W21	351	0030	04	Cso	002	В								
31 Jul S0	9W47	004	0010	01	Hsx	001	A								
01 Aug S0	8W62	006	0010	01	Axx	001	A								
02 Aug S0	9W74	004	0030	01	Cro	001	В								
03 Aug S0	9W87	004	0030	01	Axx	001	A								
								0	0	0	0	0	0	0	0
Crossed W	est Lim	b.													
Absolute h	neliograp	hic long	gitude: 351												




Recent Solar Indices (preliminary) of the observed monthly mean values

	Sunspot Numbers Radio Flux Geomagnetic												
					_		Flux	Geomagnetic					
	Observed			Smooth		*Penticton		Planetary					
Month	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value				
				,	2004								
				•	_00.								
July	87.8	51.0	0.58	68.3	40.2	118.5	105.9	23	13.8				
August	69.5	40.9	0.59	66.6	39.3	110.1	105.0	11	13.8				
September	50.0	27.7	0.55	63.7	37.6	103.1	103.7	10	13.6				
October	77.9	48.4	0.62	61.3	35.9	105.7	102.1	9	13.5				
November		43.7	0.62	60.0	35.4	113.2	101.5	26	14.1				
December	34.7	17.9	0.52	58.8	35.3	94.6	101.3	11	14.8				
				2	2005								
January	52.0	31.3	0.60	57.3	34.7	102.4	100.3	22	14.7				
February	45.4	29.1	0.64	56.4	34.0	97.3	98.5	11	14.6				
March	41.0	24.8	0.60	55.8	33.6	90.0	97.2	12	15.3				
April	41.5	24.4	0.59	52.6	31.7	85.9	95.5	12	15.7				
May	65.4	42.6	0.65	48.3	29.0	99.5	93.2	20	14.8				
June	59.8	39.6	0.66	47.9	28.9	93.7	91.9	13	13.9				
July	71.0	39.9	0.56	48.1	29.2	96.6	90.9	16	13.1				
August	65.6	36.4	0.55	45.4	27.5	90.7	89.3	16	12.2				
September		22.1	0.56	42.9	25.9	90.8	87.8	21	11.8				
Septemeen		22.1	0.00	.2.,	20.9)	07.0	-1	11.0				
October	13.0	8.5	0.65	42.6	25.5	76.7	87.4	7	11.6				
November	32.2	18.0	0.56	42.1	24.9	86.3	86.7	8	11.1				
December	62.6	41.2	0.66	40.1	23.0	90.8	85.4	7	10.4				
				:	2006								
τ.	20.0	15.4	0.55			02.0		_					
January	28.0	15.4	0.55			83.8		6					
February	5.3	4.7	0.89			76.6		6					
March	21.3	10.8	0.51			75.5		8					
April	55.2	30.2	0.55			89.0		11					
May	39.6	22.2	0.56			81.0		8					
June	37.7	13.9	0.37			80.1		8					

NOTE: All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI= 120.8, occurred April 2000. *After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 31 July 2006

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm² –sec –sr) with energies greater than 2 MeV at GOES-12 (W075).

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SEC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/m²⁾ as measured by GOES 12 (W75) and GOES 10 (W135) in two wavelength bands, .05 - .4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

