Space Weather Highlights 23 October – 29 October 2006

Solar activity was at very low levels. Only a few low level B-class flares were observed during the period.

No greater than 10 MeV proton events were observed.

The greater than 2 MeV electron flux at geosynchronous orbit was at high levels on 23 – 27 October.

The geomagnetic field ranged from quiet to active levels at middle latitudes, while high latitudes experienced quiet to major storm conditions. The period began under the tail end of a coronal hole high speed stream. Solar wind speed at ACE was decreasing from a high around 525 km/s while the IMF Bz did not vary much beyond +/- 5 nT. As a result, the geomagnetic field was at quiet to unsettled levels. Wind speed continued to decrease to a low of approximately 300 km/s midday on 27 October. Late on the 27th, density, temperature and wind speed increased, all indicative of a corotating interaction region in advance of a coronal hole high speed stream. Early on 28 October, the IMF Bz began fluctuating between +/- 10 nT and by the end of 28 October, the geomagnetic field had responded with unsettled to active conditions at middle latitudes, and active to major storm conditions at high latitudes. By the end of the summary period, wind speed increased to near 600 km/s and the geomagnetic field remained at similar storm levels.

Space Weather Outlook 01 November – 27 November 2006

Solar activity is expected to be at very low to low levels.

No greater than 10 MeV proton events are expected.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at high levels on 01 - 03 November, 10 - 23 November, and 25 - 26 November.

The geomagnetic field is expected to be mostly quiet to unsettled for the majority of the forecast period. Recurrent coronal hole high speed wind streams are expected to rotate into geoeffective positions on 09 - 11 November, 17 November, and again on 24 - 25 November. Unsettled to minor storm periods are possible on 09 - 11 November and 24 - 25 November, while unsettled to active levels are expected on 17 November.



				Daily So	ur De	aa						
	Radio	adio Sun Sunspot X-ray Flares										
	Flux	spot	Area Background		X-ray Flux				Op			
Date	10.7 cm	No.	(10 ⁻⁶ hemi.))	С	М	Х	S	1	2	3	4
23 October	76	50	210	A3.4	0	0	0	0	0	0	0	0
24 October	75	35	170	A3.0	0	0	0	0	0	0	0	0
25 October	75	15	100	A2.9	0	0	0	0	0	0	0	0
26 October	72	0	0	A2.7	0	0	0	0	0	0	0	0
27 October	72	14	20	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
28 October	75	28	170	A1.6	0	0	0	0	0	0	0	0
29 October	73	0	0	A1.5	0	0	0	0	0	0	0	0

Daily Solar Data

Daily Particle Data

	Pro	oton Fluence	Electron Fluence					
	(proto	ons/cm ² -day-si	•)	(electrons/cm ² -day-sr)				
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV				
23 October 24 October	2.3E+6 2.6E+6	1.6E+4 1.7E+4	3.5E+3 3.5E+3	1.2E+8 1.4E+8				
25 October	1.3E+6	1.6E+4	3.6E+3	1.2E+8				
26 October	1.6E+6	1.6E+4	3.6E+3	1.7E+8				
27 October	2.3E+6	1.6E+4	3.8E+3	1.3E+8				
28 October	1.4E+6	1.6E+4	3.6E+3	1.1E+7				
29 October	1.5E+6	1.7E+4	3.6E+3	1.6E+7				

Daily Geomagnetic Data

	Duny Geomagnetic Data											
	Μ	liddle Latitude]		Estimated							
	F	redericksburg		College]	Planetary						
Date	Α	K-indices	А	K-indices	Α	K-indices						
23 October	2	0-0-0-1-0-1-1-1	3	2-0-0-2-2-1-1-0	3	1-0-0-1-1-1-1-1						
24 October	3	0-0-2-1-1-1-1	4	0-0-2-1-3-0-1-1	4	0-1-2-1-1-1-1						
25 October	4	2-2-2-1-1-1-0-0	2	0-1-2-1-0-0-0-0	4	1-2-2-1-0-0-0-0						
26 October	1	0-0-1-1-1-0-0-0	3	0-0-1-2-2-1-0-0	1	0-0-1-0-1-0-0-0						
27 October	3	1-0-0-1-1-1-2-2	2	0-0-0-2-1-0-1-1	2	1-0-0-1-0-0-1-2						
28 October	12	1-2-3-3-2-3-2-4	28	1-1-4-6-4-5-2-4	14	1-2-3-4-2-3-3-4						
29 October	13	3-3-3-2-3-2-3-3	36	4-4-6-5-5-4-3-3	21	3-4-4-3-3-3-4-4						



Date & Time of Is	ssue Type of Alert or Warning	Date & Time of Event UTC
23 Oct 0903	ALERT: Electron 2MeV Integral Flux >1000pfu	23 Oct 0845
24 Oct 0923	ALERT: Electron 2MeV Integral Flux >1000pfu	24 Oct 0905
25 Oct 0834	ALERT: Geomagnetic K=4	25 Oct 0833
25 Oct 0942	ALERT: Electron 2MeV Integral Flux >1000pfu	25 Oct 0920
26 Oct 0532	ALERT: Electron 2MeV Integral Flux >1000pfu	26 Oct 0500
27 Oct 1452	ALERT: Electron 2MeV Integral Flux >1000pfu	27 Oct 0900
28 Oct 1044	WARNING: Geomagnetic K=4	28 Oct 1100 – 1600
28 Oct 1055	ALERT: Geomagnetic K=4	28 Oct 1055
28 Oct 1555	EXTENDED WARNING: Geomagnetic K=4	28 Oct 1100 – 2359
28 Oct 2301	EXTENDED WARNING: Geomagnetic K=4	28 Oct 1100 – 29/1600
29 Oct 1933	WARNING: Geomagnetic K=4	29 Oct 1933 - 2359
29 Oct 1946	ALERT: Geomagnetic K=4	29 Oct 1940
29 Oct 2337	EXTENDED WARNING: Geomagnetic K=4	29 Oct 1933 – 30/1600



Twenty-seven Day Outlook



	Radio Flux	Planetary	Largest		Radio Fluz	x Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
01 Nov	80	5	2	15 Nov	75	5	2
02	85	5	2	16	75	10	3
03	85	8	3	17	75	15	4
04	85	8	3	18	75	10	3
05	85	5	2	19	75	5	2
06	85	5	2	20	75	5	2
07	85	5	2	21	75	5	2
08	85	5	2	22	75	5	2
09	85	20	5	23	75	8	3
10	85	15	4	24	75	12	4
11	80	12	3	25	75	20	5
12	75	10	3	26	80	8	3
13	75	8	3	27	80	5	2
14	75	5	2				



					j	Energet	ic Events					
	Time					X-ray Optical Information					Sweep Free	
Date			1/2]	Integ	Imp/	Location	Rgn	Radi	o Flux	Intensity	
	Begin	Max	Max	Class	Flux	Brtns	Lat CMD	#	245	2695	II IV	
No Even	ts Obs	erved										
						Flar	e List					
						C	Optical					
			Time	•		2	X-ray	Imp /	Lo	cation	Rgn	
Date		Begin	Max	E	ind	(Class.	Brtns	Lat	CMD	_	
23 Octobe	r	0003	000	7 0	014	I	31.9					
24 Octobe	r	No Fla	ares Ob	served								
25 Octobe	r	1859	190	6 1	913]	B1.3					
		1930	193	7 1	943]	B1.3					
26 Octobe	r	No Fla	ares Ob	served								
27 Octobe	r	No Fla	ares Ob	served								
28 Octobe	r	0526	053	2 0.	534]	B1.4					
		0654	065	8 0	700]	B1.2					
		1920	192	4 19	930	1	B1.0					
		2109	211	3 2	115	1	32.5					
29 Octobe	r	0436	044	$\frac{1}{1}$ $\frac{1}{0}$	450	1	B1.2					
	-	0.00	0.1	- 0								



				Reg	gion Su	mmary	y									
	Locatio	n		Sunspot	Character Flares	istics										
		Helio	Area	Extent	Spot	Spot	Mag		X-ra	y		(Optic	al		
Date	(°Lat°CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	М	Х	S	1	2	3	4	
	Re	gion 917	7													
19 Oct	S05W04	344	0030	04	Bxo	004	В									
20 Oct	S05W20	347	0030	04	Bxo	005	В				1					
21 Oct	S05W33	347	0070	04	Cso	006	В									
22 Oct	S05W46	346	0200	08	Dac	008	Bg									
23 Oct	S05W60	347	0160	10	Dai	024	Bg									
24 Oct	S05W73	347	0140	11	Cao	013	B									
25 Oct	S05W86	347	0100	10	Cao	005	В									
								0	0	0	1	0	0	0	0	
Crosse	d West Lim	ıb.														
Absolu	ite heliogra	ohic long	gitude: 344													
	Re	gion 918	3													
23 Oct	S04W72	359	0050	04	Cro	006	В									
24 Oct	S04W85	359	0030	02	Hrx	002	А									
								0	0	0	0	0	0	0	0	
Crosse	d West Lim	ıb.														
Absolu	ite heliogra	ohic long	gitude: 359													
	Re	gion 919)													
27 Oct	S15W01	236	0020	04	Bvo	004	B									
27 Oct	S15W01	230	0020	04	Bxo	004	B									
20 Oct	S15W15	236	0020	04	DAO	005	D									
27 000	515 1 20	230						0	0	0	0	0	0	0	0	
Still or	Dick							0	0	0	0	0	U	U	0	
Absolu	i Disk. ite helioarai	phic long	ritude 236													
Auson	ne nenograj		situae. 250													
	Re	gion 920)													
28 Oct	S07W82	303	0150	12	Eso	005	В	0	0	0	0	0	0	0	0	
Still or	n Diek							U	U	U	U	U	U	0	0	
Absolu	ite helioara	hic lone	ritude: 302													
AUSUI	ne nenograj		51100C. 303													_



Sunspot Numbers Radio Flux Geomagnetic											
(Theoryad	values	Ratio	Smooth	values	*Penticton	Smooth	Planetary	Smooth		
Month	SEC		RI/SEC	SEC	PI	10.7 cm	Value	<u>Γ Tanctal y</u> Δ n	Value		
wionui	JEC	<u>IXI</u>	MOLU	JLC	1/1	10.7 011	v aluc	- Ар	v and		
					2004						
		10.1	0.60	<i>(</i> 1 0	25.0	1055	100.1	0	10 5		
October	77.9	48.4	0.62	61.3	35.9	105.7	102.1	9	13.5		
November	70.5	43.7	0.62	60.0	35.4	113.2	101.5	26	14.1		
December	34.7	17.9	0.52	58.8	35.3	94.6	101.3	11	14.8		
					2005						
January	52.0	31.3	0.60	57.3	34.7	102.4	100.3	22	14.7		
February	45.4	29.1	0.64	56.4	34.0	97.3	98.5	11	14.6		
March	41.0	24.8	0.60	55.8	33.6	90.0	97.2	12	15.3		
April	<i>A</i> 1 5	24.4	0 59	52.6	31.7	85.9	95 5	12	157		
May	-1.J 65 /	<u>2</u> т.т Л2 б	0.55	<i>J</i> 2.0 <i>I</i> 8.3	20.0	00.5	03.2	$\frac{12}{20}$	1/1 8		
June	50.8	42.0 30.6	0.05	40.5	29.0 28.0	03.7	01.0	13	13.0		
June	39.0	39.0	0.00	47.7	20.9	95.7	91.9	15	13.9		
July	71.0	39.9	0.56	48.1	29.2	96.6	90.9	16	13.1		
August	65.6	36.4	0.55	45.4	27.5	90.7	89.3	16	12.2		
September	39.2	22.1	0.56	42.9	25.9	90.8	87.8	21	11.8		
October	13.0	8.5	0.65	42.6	25.5	76.7	87.4	7	11.6		
November	32.2	18.0	0.56	42.1	24.9	86.3	86.7	8	11.1		
December	62.6	41.2	0.66	40.1	23.0	90.8	85.4	7	10.4		
					2006						
January	28.0	15.4	0.55	37.2	20.8	83.8	84.0	6	9.9		
February	5.3	4.7	0.89	33.4	18.7	76.6	82.6	6	9.2		
March	21.3	10.8	0.51	31.0	17.4	75.5	81.6	8	8.4		
1	55.0	20.0	0.55			00.0		11			
April	55.2 20.6	30.2	0.55			89.0		11			
May	39.6 07.7	22.2	0.56			81.0		8			
June	31.1	13.9	0.37			80.1		8			
Julv	22.6	12.2	0.54			75.8		7			
August	22.8	12.9	0.57			79.0		9			
September	25.2	14.5	0.58			77.8		8			

Recent Solar Indices (preliminary) of the observed monthly mean values

<u>NOTE</u>: All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI = 120.8, occurred April 2000. *After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 23 October 2006

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.

Electrons plot contains the five-minute averaged integral electron flux (electrons/cm²-sec -sr) with energies greater than 2 MeV at GOES-12 (W075).

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SEC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/ m^{2}) as measured by GOES 12 (W075) and GOES 11 (W135) in two wavelength bands, .05 - . 4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

