### Space Weather Highlights 23 – 29 June 2008

SEC PRF 1713 01 July 2008

Solar activity was very low. No flares were detected. The visible disk was spotless.

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit reached high levels during 23 - 24, 27, and 29 June.

Geomagnetic field activity was at quiet levels until late on 24 June when activity increased to unsettled levels. A further increase to active levels occurred during 25 - 26 June with minor to major storm levels at high latitudes. Activity decreased to quiet to unsettled levels for the remainder of the period. ACE solar wind measurements indicated a co-rotating interaction region (CIR) occurred during 24 - 25 June in advance of a recurrent coronal hole high-speed stream (HSS). The CIR was associated with increased proton densities (peak 20 p/cc at 24/2251 UTC), a solar sector boundary crossing (toward (-) to away (+)), and changes in the interplanetary magnetic field (IMF). IMF changes included increased Bt (peak 15 nT at 25/1710 UTC) and intermittent periods of southward Bz (minimum -10 nT at 25/1735 UTC). The HSS commenced on 25 June, reached a peak velocity of 686 km/sec at 26/1846 UTC, then began to gradually subside on 28 June. Solar wind velocities decreased to a low of 470 km/sec by the end of the period.

# Space Weather Outlook 02 – 28 July 2008

Solar activity is expected to be very low.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to reach high levels during 13 - 21 July.

Geomagnetic field activity is expected to be at quiet to unsettled levels during 02 - 11 July. Activity is expected to increase to active levels during 12 - 13 July with storm levels possible at high latitudes due to a recurrent coronal hole high-speed stream. Activity is expected to decrease to unsettled levels during 14 - 17 July as the high-speed stream gradually subsides. Activity is expected to decrease to quiet levels during 18 - 21 July. Activity is expected to increase to unsettled levels during 22 - 24 July due to recurrence. Activity is expected to decrease to quiet levels during 25 - 28 July.



Daily Solar Data

				Dully 50	<i></i>	an						
	Radio	Sun	Sunspot	X-ray	_			Flares				
	Flux	spot	Area	Background	X	ray F	lux		Op	otical		
Date	10.7 cm	No.	(10 <sup>-6</sup> hemi.)	)	С	M	X	S	1	2	3	4
23 June	65	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
24 June	66	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
25 June	66	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
26 June	65	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
27 June	66	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
28 June	66	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
29 June	67	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0

### Daily Particle Data

		oton Fluence	,	Electron Fluence				
	(proto	ons/cm <sup>2</sup> -day-si	r)	(electrons/cm <sup>2</sup> -day-sr)				
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV				
23 June	6.0E+5	1.8E+4	4.3E+3	7.3E+7				
24 June	9.9E+5	1.8E+4	4.3E+3	8.3E+7				
25 June	4.3E+5	1.8E+4	4.7E+3	1.4E+6				
26 June	1.2E+6	1.7E + 4	3.9E+3	4.8E+6				
27 June	1.2E+6	1.8E + 4	4.1E+3	3.6E+7				
28 June	1.0E+6	1.7E+4	4.2E+3	3.8E+7				
29 June	1.1E+6	1.8E+4	4.1E+3	4.2E+7				

Daily Geomagnetic Data

Daily Geomagnetic Data												
	M	Middle Latitude		High Latitude	E	Estimated						
	F	redericksburg		College	I	Planetary						
Date	A	K-indices	Α	K-indices	A	K-indices						
23 June	2	1-2-0-0-0-1-0	4	1-1-1-3-2-0-0-0	3	1-2-0-0-1-1-0-1						
24 June	3	0-0-0-0-1-2-3	2	1-1-0-0-0-1-1-2	4	1-1-0-0-0-1-2-3						
25 June	10	3-3-3-1-1-2-2-3	10	2-3-3-2-1-2-3-2	12	4-3-3-1-1-2-3-3						
26 June	17	4-4-2-4-4-2-2-2	23	3-3-3-6-5-1-2-1	16	4-4-2-4-3-2-3-2						
27 June	7	3-2-1-2-2-2-1	16	3-2-2-5-4-3-1-2	8	3-2-2-2-2-2						
28 June	6	1-2-1-1-1-3-2	6	1-2-1-2-2-3-1-1	7	2-2-1-2-1-2-3-2						
29 June	7	2-2-1-2-2-1-2-3	8	2-3-1-3-2-1-2-2	8	2-3-1-2-1-1-2-3						




#### Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
23 Jun 1237	ALERT: Electron 2MeV Integral Flux >1000pfu	23 Jun 1215
24 Jun 1201	ALERT: Electron 2MeV Integral Flux >1000pfu	24 Jun 1140
25 Jun 0430	WARNING: Geomagnetic K=4	25 Jun 0431 – 1600
25 Jun 0542	ALERT: Geomagnetic K=4	25 Jun 0541
25 Jun 1928	WARNING: Geomagnetic K=4	25 Jun 1930 - 2359
26 Jun 0242	WARNING: Geomagnetic K=4	26 Jun 0242 - 0700
26 Jun 0246	ALERT: Geomagnetic K=4	26 Jun 0244
26 Jun 0438	ALERT: Geomagnetic K=4	26 Jun 0435
26 Jun 0916	WARNING: Geomagnetic K=4	26 Jun 0915 - 1600
26 Jun 0919	ALERT: Geomagnetic K=4	26 Jun 0918
27 Jun 0225	WARNING: Geomagnetic K=4	27 Jun 0226 - 0700
27 Jun 0245	ALERT: Geomagnetic K=4	27 Jun 0245
27 Jun 1421	ALERT: Electron 2MeV Integral Flux >1000pfu	27 Jun 1405
29 Jun 1421	ALERT: Electron 2MeV Integral Flux >1000pfu	29 Jun 1405



#### Twenty-seven Day Outlook



Largest Daily Kp Index

	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
02 Jul	66	5	2	16 Jul	66	8	3
03	66	8	3	17	66	10	3
04	66	12	3	18	66	5	2
05	66	10	3	19	66	5	2
06	66	5	2	20	66	5	2
07	66	5	2	21	66	5	2
08	66	5	2	22	66	10	3
09	66	5	2	23	66	10	3
10	66	5	2	24	66	8	3
11	66	10	3	25	66	5	2
12	66	15	4	26	66	5	2
13	66	15	4	27	66	5	2
14	66	10	3	28	66	5	2
15	66	10	3				



**Energetic Events** 

				Ditti gu	ic Dienis				
	Time		X-ray	Opt	ical Information	1	Peak	Sweep Freq	
Date		1/2	Integ Imp/		Location	Rgn	Radio Flux	Intensity	
	Begin Max	Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV	
No E	Events Observed								

Flare List

			r iiire List					
Ti	ime		Optical X-ray	Imp /	Location	Rgn		
Begin M	1ax	End	Class.	Brtns	Lat CMD			
No Flares	Observe	d						
No Flares	No Flares Observed							
No Flares	No Flares Observed							
No Flares	Observe	d						
No Flares	Observe	d						
No Flares	Observe	d						
No Flares	Observe	d						
	Begin No Flares	No Flares Observe No Flares Observe No Flares Observe No Flares Observe No Flares Observe	Begin Max End  No Flares Observed  No Flares Observed	Time X-ray Begin Max End Class.  No Flares Observed	Time X-ray Imp/ Begin Max End Class. Brtns  No Flares Observed	Time X-ray Imp / Location Begin Max End Class. Brtns Lat CMD  No Flares Observed		

Region Summary

	Location		Sunspot	Characte	ristics					
				Flares						
	Helio	Area	Extent	Spot	Spot	Mag	X-ray		Optical	_
Date	(° Lat ° CMD) Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C M X	S 1	2 3	4

#### Region 999

16 Jun S02E60	204	0020	01	Axx	001	A
17 Jun S02E46	205	0020	01	Hsx	001	A
18 Jun S03E37	204	0010	01	Hsx	001	A
19 Jun S03E20	205	0010	01	Axx	001	A
20 Jun S02E06	206	0010	01	Hsx	001	A
21 Jun S02W08	207	0010	01	Hsx	001	A
22 Jun S02W22	207	0010	01	Hsx	001	A
22 Ivn C02W25	207					

23 Jun S02W35 207

24 Jun S02W48 207

25 Jun S02W61 207

26 Jun S02W74 206

27 Jun S02W87 206

0 0 0 0 0 0 0 0

Crossed West Limb.

Absolute heliographic longitude: 206




# Recent Solar Indices (preliminary) Of the observed monthly mean values

	Sunsp					Flux	Geoma	gnetic				
Observed				values	*Penticton	Smooth	Planetary	Smooth				
SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value		Value				
			,	2006			-					
22.8	12.9	0.57	27.6	15.6	79.0	80.3	9	8.7				
25.2	14.5	0.58	27.7	15.6	77.8	80.2	8	8.7				
15.7	10.4	0.66	25.2	14.2	74.3	79.4	8	8.6				
31.5	21.5	0.68	22.3	12.7	86.4	78.5	9	8.5				
22.2	13.6	0.61	20.7	12.1	84.3	77.9	15	8.5				
<b>2007</b> January 26.6 16.9 0.64 19.7 12.0 83.5 77.5 6 8.4												
								8.4				
								8.4				
9.7	4.8	0.49	17.5	10.8	72.3	76.0	8	8.4				
								8.5				
								8.4				
20.0	12.0	0.60	12.8	7.7	73.7	73.2	7	7.8				
15.6	10.0	0.64	11.6	7.0	71.6	72.5	8	7.4				
								7.6				
								7.8				
		<b>0.0</b> 0	J.,		0,,1	, 1,0		7.0				
1.3	0.9	0.70	10.0	6.1	65.5	71.5	9	7.9				
2.5	1.7	0.68	9.4	5.7	69.7	71.1	5	7.8				
16.2	10.1	0.62			78.6		4					
				2008								
15.9	9.3	0.58			72.9		10					
49	2.9	0.59			70.3		9					
5.7	2.9	0.51			68.4		6					
	22.8 25.2 15.7 31.5 22.2 26.6 17.2 9.7 6.9 19.4 20.0 15.6 9.9 4.8 1.3 2.5 16.2 5.1 3.8 15.9	Observed values           SEC         RI           22.8         12.9           25.2         14.5           15.7         10.4           31.5         21.5           22.2         13.6           26.6         16.9           17.2         10.6           9.7         4.8           6.9         3.7           19.4         11.7           20.0         12.0           15.6         10.0           9.9         6.2           4.8         2.4           1.3         0.9           2.5         1.7           16.2         10.1           5.1         3.4           3.8         2.1           15.9         9.3           4.9         2.9	Sunspot Numbe           Observed values         Ratio           SEC         RI         RI/SEC           22.8         12.9         0.57           25.2         14.5         0.58           15.7         10.4         0.66           31.5         21.5         0.68           22.2         13.6         0.61           26.6         16.9         0.64           17.2         10.6         0.62           9.7         4.8         0.49           6.9         3.7         0.54           19.4         11.7         0.60           20.0         12.0         0.60           15.6         10.0         0.64           9.9         6.2         0.63           4.8         2.4         0.50           1.3         0.9         0.70           2.5         1.7         0.68           16.2         10.1         0.62           5.1         3.4         0.67           3.8         2.1         0.55           15.9         9.3         0.58           4.9         2.9         0.59	Sunspot Numbers           Observed values         Ratio Ratio SEC         Smooth SEC           22.8         12.9         0.57         27.6           25.2         14.5         0.58         27.7           15.7         10.4         0.66         25.2           31.5         21.5         0.68         22.3           22.2         13.6         0.61         20.7           26.6         16.9         0.64         19.7           17.2         10.6         0.62         18.9           9.7         4.8         0.49         17.5           6.9         3.7         0.54         16.0           19.4         11.7         0.60         14.2           20.0         12.0         0.60         12.8           15.6         10.0         0.64         11.6           9.9         6.2         0.63         10.2           4.8         2.4         0.50         9.9           1.3         0.9         0.70         10.0           2.5         1.7         0.68         9.4           16.2         10.1         0.62           5.1         3.4         0.67	Sunspot Numbers   Smooth values   SEC   RI   RI/SEC   SEC   RI	Observed values SEC         RI         Ratio RI/SEC         Smooth values SEC         *Penticton 10.7 cm           22.8         12.9         0.57         27.6         15.6         79.0           25.2         14.5         0.58         27.7         15.6         77.8           15.7         10.4         0.66         25.2         14.2         74.3           31.5         21.5         0.68         22.3         12.7         86.4           22.2         13.6         0.61         20.7         12.1         84.3           2007           26.6         16.9         0.64         19.7         12.0         83.5           17.2         10.6         0.62         18.9         11.6         77.8           9.7         4.8         0.49         17.5         10.8         72.3           6.9         3.7         0.54         16.0         9.9         72.4           19.4         11.7         0.60         14.2         8.7         74.5           20.0         12.0         0.60         12.8         7.7         73.7           15.6         10.0         0.64         11.6         7.0         71.6	Sunspot Numbers         Radio Flux           Observed values         Ratio RI/SEC         SEC         RI         *Penticton         Smooth Smooth 10.7 cm         Value           2006           22.8         12.9         0.57         27.6         15.6         79.0         80.3           25.2         14.5         0.58         27.7         15.6         77.8         80.2           15.7         10.4         0.66         25.2         14.2         74.3         79.4           31.5         21.5         0.68         22.3         12.7         86.4         78.5           22.2         13.6         0.61         20.7         12.1         84.3         77.9           26.6         16.9         0.64         19.7         12.0         83.5         77.5           17.2         10.6         0.62         18.9         11.6         77.8         76.9           9.7         4.8         0.49         17.5         10.8         72.3         76.0           6.9         3.7         0.54         16.0         9.9         72.4         75.2           19.4         11.7         0.60         14.2         8	Sunspot Numbers				

**NOTE:** All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI= 120.8, occurred April 2000. \*After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 23 June 2008

Protons plot contains the five-minute averaged integral proton flux (protons/cm<sup>2</sup>-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm<sup>2</sup>-sec -sr) with energies greater than 2 MeV at GOES-12 (W075).

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SEC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





#### Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/ $m^2$ ) as measured by GOES 10 (W060) and GOES 11 (W135) in two wavelength bands, .05 - .4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm<sup>2</sup> –sec-sr) as measured by GOES-11 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm<sup>2</sup>-sec-sr) at greater than 10 MeV.



## M5 or Greater X-Ray Flares

