Space Weather Highlights 31 August – 06 September 2009

SWO PRF 1775 08 September 2009

Solar activity was very low. Region 1025 (N17, L=170, class/area Bxo/010 on 01 September) was numbered on 31 August. Region 1025 remained inactive and decayed to spotless plage on 02 September.

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit was at normal levels during the period.

Geomagnetic field activity was at predominantly quiet levels during the period. Unsettled to active periods were observed at high latitudes between 04/0000 UTC to 04/0900 UTC. Solar wind data at ACE indicated the increased activity was due to a recurrent coronal hole high speed stream (CH HSS). Velocities at ACE increased from a low of 342 km/s at 03/1259 UTC to a high of 504 km/s at 04/1831 UTC, before gradually decreasing to 371 km/s at the end of the period. Interplanetary magnetic field activity included an increase in Bt between 03/1344 UTC and 04/1208 UTC (maximum of 10 nT at 04/0150), while Bz varied between -/+ 6 nT. Density reached a maximum of 10 p/cc at 04/0528 UTC.

Space Weather Outlook 09 September – 05 October 2009

Solar activity is expected to be very low.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at normal to moderate flux levels during the period.

Geomagnetic field activity is expected to be mostly quiet, with isolated unsettled levels during 09 – 10 September due to a weak CH HSS. Quiet levels are expected during 11 – 14 September. Quiet to unsettled conditions, with isolated active levels are expected during 15 – 17 September due to a CH HSS. Predominantly quiet levels are expected during 18 – 25 September. Quiet to unsettled conditions, with isolated active levels are expected on 26 September due to a CH HSS. Activity is expected to decrease to predominantly quiet levels during 27 September – 05 October.



Daily Solar Data

	Radio	Sun	Sunspot	X-ray	_		Flares								
	Flux	spot	Area	Area Background		ray Fl	lux		Optical						
Date	10.7 cm	No.	(10 ⁻⁶ hemi.)	С	M	X	S	1	2	3	4			
31 August	68	12	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
01 Septemb	er 69	12	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
02 Septemb	er 68	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
03 Septemb	er 69	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
04 Septemb	er 68	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
05 Septemb	er 69	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			
06 Septemb	er 69	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0			

Daily Particle Data

		on Fluence	Electron Fluence						
	(proton	s/cm ² -day-sr)	(electrons/cm ² -day-sr)						
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV					
31 August	5.8e + 05	1.9e + 04	4.1e+03	2.3e+05					
01 September	7.2e+05	1.9e + 04	4.2e+03	8.2e + 05					
02 September	7.1e+05	2.0e+04	4.3e+03	7.7e+05					
03 September	1.2e+06	2.1e+04	4.4e+03	1.2e + 06					
04 September	4.7e+05	1.9e + 04	4.2e+03	1.9e + 05					
05 September	6.6e+05	1.9e + 04	4.2e+03	2.7e + 05					
06 September	8.2e+05	1.9e+04	4.4e+03	3.4e+05					

Daily Geomagnetic Data


	N	Middle Latitude		High Latitude		Estimated
	I	Fredericksburg		College]	Planetary
Date	Α	K-indices	A	K-indices	A	K-indices
31 August	5	2-1-1-2-2-1-1-1	4	2-1-1-2-2-2-1-0	5	2-1-1-1-2-2-1-1
01 September	2	1-1-0-0-2-0-0-0	2	1-0-0-2-2-0-0-0	4	1-0-0-1-1-2-1-1
02 September	2	2-1-0-0-1-1-0-1	0	0-1-0-0-0-0-0	3	2-1-0-0-1-0-1-1
03 September	4	0-0-1-1-1-2-2-2	2	1-0-1-1-0-1-1-1	4	1-0-1-1-1-2-1-2
04 September	4	2-2-1-2-1-1-0	9	2-3-3-4-1-1-0-0	5	2-2-1-2-1-1-1
05 September	2	1-1-1-1-0-0-1	0	0-1-0-0-0-0-0	2	1-2-0-0-0-0-1
06 September	2	0-2-0-1-0-1-1-0	1	0-1-0-0-0-0-1	3	0-2-0-1-0-0-1-1

Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
No Alorts Issued		



Twenty-seven Day Outlook



Largest Daily Kp Index

	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
09 Sep	69	10	3	23 Sep	68	5	2
10	69	8	3	24	68	5	2
11	69	5	2	25	68	5	2
12	68	5	2	26	68	10	4
13	68	5	2	27	68	5	2
14	68	5	2	28	69	5	2
15	68	8	3	29	69	5	2
16	68	8	3	30	69	5	2
17	68	5	2	01 Oct	69	5	2
18	68	5	2	02	69	5	2
19	68	5	2	03	69	5	2
20	68	5	2	04	69	5	2
21	68	5	2	05	69	8	3
22	68	5	2				



Energ	etic	Events
Livers	cuc	LIVETUS

Time		X-ray Optical Information				Peak	Sweep Freq				
Date	Date		1/2	Integ	Imp/	Location	Rgn	Radio Flux	Intensity		
	Begin	Max	Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV		
No Event	s Observ	ed									

Flare List

				Ture List			
	Time			X-ray	Imp/	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	
31 August	No Fl	ares Obs	erved				
01 September	No Fl	ares Obs	erved				
02 September	No Fl	ares Obs	erved				
03 September	No Fl	ares Obs	erved				
04 September	No Fl	ares Obs	erved				
05 September	No Fl	ares Obs	erved				
06 September	No Fl	ares Obs	erved				

Region Summary

	Location			Sunspot Characteristics						Flares							
	Helio		Area	Area Extent		Spot Mag			X-ray		Optical						
Date	(° Lat ° CMD) Lon (10°		(10 ⁻⁶ hemi)	emi) (helio) Class Cou			Class	(C	M	X	S	1	2	3	4	
	$R \rho$	gion 10	025														

31 Aug	N17E32	169	10	3	BXO	2	В
01 Sep	N17E18	170	10	3	BXO	2	В

01 Sep N17E18 170 10 3 BXO 2 ... 02 Sep N17E05 170

03 Sep N17W08 170 04 Sep N17W21 170

06 Sep N17W47 170 0 0 0 0 0 0 0 0 0

Still on Disk.

05 Sep N17W34 170

Absolute heliographic longitude: 170




Recent Solar Indices (preliminary) Of the observed monthly mean values

		Sunsn	ot Numbe		···········y	Radio	Flux	Geoma	gnetic
	Observed	_		Smooth	values	*Penticton		Planetary	-
Month	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value
					007			•	
September	4.8	2.4	0.50	9.9	5.9	67.1	71.5	9	7.8
-									
October	1.3	0.9	0.70	10.0	6.0	65.5	71.5	9	7.9
November		1.7	0.68	9.4	5.7	69.7	71.1	5	7.8
December	16.2	10.1	0.62	8.1	4.9	78.6	70.5	4	7.8
				2	000				
Ionuomi	5 1	3.3	0.67	6.9	008 4.2	72.1	70.0	6	7.7
January February	5.1 3.8	2.1	0.67	5.9	3.6	71.2	69.6	6 9	7.7 7.6
March	3.8 15.9	9.3	0.58	5.3	3.3	72.9	69.5	10	7.0 7.4
Maich	13.9	9.3	0.56	5.5	3.3	12.9	09.3	10	/ . 4
April	4.9	2.9	0.59	5.3	3.4	70.3	69.6	9	7.1
May	5.7	3.2	0.51	5.7	3.5	68.4	69.7	6	6.9
June	4.2	3.4	0.74	5.2	3.3	65.9	69.2	7	6.8
July	1.0	0.8	0.50	4.5	2.8	65.8	68.8	6	6.6
August	0.0	0.5	-1.00	4.4	2.7	66.4	68.6	5	6.2
September	1.5	1.1	0.73	3.7	2.3	67.1	68.4	5	5.8
October	5.2	2.9	0.56	2.9	1.9	68.3	68.2	6	5.4
November		4.1	0.60	2.7	1.8	68.6	68.3	3	5.1
December		0.8	0.62	2.7	1.7	69.2	68.5	2	4.9
_					009			_	
January	2.8	1.3	0.54	3.0	1.8	69.8	68.7	3	4.8
February	2.5	1.4	0.56	3.1	1.9	70.0	68.9	4	4.7
March	0.7	0.7	1.00			69.2		4	
April	1.2	1.2	1.00			69.7		4	
May	3.9	2.9	0.74			70.6		4	
June	6.6	2.6	0.39			68.6		5	
July	5.0	3.5	0.70			68.3		5	
August	0.3	0.0	0.00			67.3		5	
0	- · -							-	

NOTE: All values are preliminary estimates except for RI Sunspot Numbers which are final values when available.

Cycle 23 started in May 1996 with an RI=8.0. Cycle 23 maximum was April 2000 with an RI=120.8. **SWPC sunspot number was less than RI value, so a ratio could not be computed.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 31 August 2009

GOES-11 designated Primary Electron Satellite and GOES-10 Secondary: December 1, 2008 the GOES-12 Electron sensor began experiencing periods of noise and sensor is unreliable.

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm²-sec -sr) with energies greater than 2 MeV at GOES-11.

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-11. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWPC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/ m^2) as measured by GOES 10 (W060) and GOES 11 (W135) in two wavelength bands, .05 - .4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

