#### Space Weather Highlights 02 November – 08 November 2009

SWO PRF 1784 03 November 2009

Solar activity was very low. Region 1030 (N25, L=046, class/area Cro/050 on 05 November) emerged on the disk on 05 November. Region 1030 gradually decayed to spotless plage late on 07 November.

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit was at normal levels during the period.

The geomagnetic field was at quiet levels through most of the period. On 08 November, a weak recurrent coronal hole high-speed stream (CH-HSS) was observed from the ACE spacecraft. Quiet to isolated unsettled levels were observed between 08/0900-2100 UTC, with isolated active levels at high latitudes between 08/1200-1800 UTC. Velocities increased from 247 km/s at 07/1331 UTC to a high of 461 km/s at 08/2145 UTC. Interplanetary magnetic field activity associated with the weak CH HSS included an increase in Bt (maximum of 11 nT at 08/0957 UTC) and southward Bz (minimum of -9 nT at 08/1821Z).

### Space Weather Outlook 11 November 2009 – 07 December 2009

Solar activity is expected to be very low to low. There may be a small increase in background levels from 15-27 November due to the return of Region 1029.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at normal levels through the period.

The geomagnetic field is expected to be at quiet levels during 11-17 November. Quiet to unsettled periods, with isolated active levels, are expected on 18-20 November, due to possible effects from a recurrent CH-HSS. Activity is expected to decrease to quiet levels on 21 November for the remainder of the period.



## Daily Solar Data

	Radio	Sun	Sunspot X-ray		Flares							
	Flux	spot	Area	Background	X-	-ray F	lux		O	ptical		
Date	10.7 cm	No.	(10 <sup>-6</sup> hemi.	.)	С	M	X	S	1	2	3	4
02 Novemb	er 71	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
03 Novemb	er 72	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
04 Novemb	er 71	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
05 Novemb	er 71	15	50	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
06 Novemb	er 71	16	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
07 Novemb	er 71	11	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
08 Novemb	er 71	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0

## Daily Particle Data

		on Fluence as/cm <sup>2</sup> -day-sr)	Electron Fluence (electrons/cm²-day-sr)				
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV			
02 November	8.3e+05	2.0e+04	4.5e+03	1.2e+06			
03 November	6.2e+05	2.0e+04	4.3e+03	5.7e+05			
04 November	1.1e+06	1.9e + 04	4.6e + 03	1.0e+06			
05 November	1.1e+06	1.9e + 04	4.2e+03	1.1e+06			
06 November	8.9e+05	2.0e+04	4.4e+03	8.7e+05			
07 November	9.7e+05	2.0e+04	4.4e+03	1.1e+06			
08 November	8.0e+05	2.0e+04	4.4e+03	3.5e+05			


Daily Geomagnetic Data

	Middle Latitude			High Latitude	]	Estimated
		Fredericksburg		College	]	Planetary
Date	Α	K-indices	Α	K-indices	Α	K-indices
02 November	2	1-1-2-1-0-1-0-0	1	0-0-2-1-0-0-0	2	0-0-1-1-0-0-0-0
03 November	1	0-0-0-0-1-1-1-0	0	0-0-0-0-0-0-0	0	0-0-0-0-0-0-0
04 November	1	0-0-0-0-1-0-1-0	0	0-0-0-0-0-0-0	0	0-0-0-0-0-0-0
05 November	0	0-0-1-0-0-0-0	0	0-0-1-0-0-0-0	1	0-0-1-0-0-0-0
06 November	0	0-0-0-0-0-0-0	0	0-0-0-0-0-0-0	0	0-0-0-0-0-0-0
07 November	0	0-0-0-0-0-0-0	0	0-0-0-0-0-0-0	1	0-0-0-0-1-0-1
08 November	4	0-1-1-2-2-2-1-1	11	1-0-0-3-4-4-2-2	6	0-1-1-1-2-2-3-2

Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
No Alerts Issued		





	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
11 Nov	72	5	2	25 Nov	75	5	3
12	72	5	2	26	75	8	4
13	72	5	2	27	72	5	2
14	73	5	2	28	72	5	2
15	73	5	2	29	70	5	2
16	73	5	2	30	70	5	2
17	73	5	2	01 Dec	70	5	2
18	74	10	4	02	70	5	2
19	75	8	4	03	70	5	2
20	75	8	4	04	70	5	2
21	75	5	3	05	70	5	2
22	80	5	2	06	70	5	2
23	80	5	2	07	72	5	2
24	80	5	2				



Energetic Events

				-	Bitter get	10 23 / 0.1115			
	Time			X-ray	Optical Information			Peak	Sweep Freq
Date	'		1/2	Integ	Imp/	Location	Rgn	Radio Flux	Intensity
	Begin	Max	Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV
No Event	s Observ	red							

#### Flare List

					Optical	
	Time		X-ray	Imp/	Location	Rgn
Date	Begin Max	End	Class.	Brtns	Lat CMD	
02 November	No Flares Obs	served				
03 November	No Flares Obs	served				
04 November	No Flares Obs	served				
05 November	No Flares Obs	served				
06 November	No Flares Obs	served				
07 November	No Flares Obs	served				
08 November	No Flares Obs	served				

## Region Summary

	Locatio		Sunspot	t Characte	ristics	Flares							
		Helio	Area	Extent	Spot	Spot	Mag	X-ray Optical					
Date	(° Lat ° CMD)	Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C M X S 1 2 3 4					
	Region 1030												
05 N	ov N25E04	46	50	6	CRO	5	В						
06 N	ov N24W07	7 44	10	5	BXO	6	В						
07 N	ov N25W25	5 48	10	1	AXX	1	A						
08 N	ov N25W38	8 48											
								$0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0$					
Still	on Disk.												
Abso	lute heliogra	phic lo	ngitude: 46										




Recent Solar Indices (preliminary)
Of the observed monthly mean values

	ot Number	Radio	Flux	Geomagnetic					
	Observed	_		Smooth	values	*Penticton		Planetary	
Month	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value
November	2.5	1.7	0.68	9.4	5.7	69.6	71.4	7	7.8
December	16.2	10.1	0.62	8.1	4.9	78.5	70.8	6	7.8
				2	008				
January	5.1	3.3	0.65	6.9	4.2	74.3	70.3	8	7.8
February	3.8	2.1	0.55	5.9	3.6	71.1	69.9	11	7.6
March	15.9	9.3	0.58	5.3	3.3	72.9	69.8	11	7.5
April	4.9	2.9	0.59	5.3	3.4	70.2	69.8	9	7.3
May	5.7	3.2	0.56	5.7	3.5	68.4	69.8	6	7.2
June	4.2	3.4	0.81	5.2	3.3	65.9	69.4	7	7.0
								_	- 0
July	1.0	0.8	0.80	4.5	2.8	65.7	68.8	5	6.8
August	0.0	0.5	**	4.4	2.7	66.3	68.6	5	6.3
September	1.5	1.1	0.73	3.7	2.3	67.1	68.4	6	5.8
Ootobor	5.2	2.0	0.56	2.0	1.0	60.2	68.2	7	5 1
October November	5.2	2.9	0.56	2.9	1.9	68.3			5.4
		4.1	0.60	2.7	1.8	68.6	68.3	4	5.1
December	1.3	0.8	0.62	2.7	1.7	69.2	68.5	4	4.9
				2	009				
January	2.8	1.3	0.46	3.0	1.8	69.8	68.7	4	4.7
February	2.5	1.4	0.56	3.1	1.9	70.0	68.8	5	4.7
March	0.7	0.7	1.00	3.4	2.0	69.2	69.0	5	4.6
April	1.2	1.2	1.00			69.7		4	
May	3.9	2.9	0.74			70.5		4	
June	6.6	2.6	0.39			68.6		4	
July	5.0	3.5	0.70			68.2		4	
August	0.3	0.0	0.00			67.4		5	
September	6.6	4.2	0.64			70.5		4	
October	7.0	4.6	0.66			72.4		3	

**NOTE:** Values are final except for the most recent 6 months which are considered preliminary. Cycle 23 started in May 1996 with an RI=8.0. Cycle 23 maximum was April 2000 with an RI=120.8.



<sup>\*\*</sup> SWPC sunspot number was less than RI value, so a ratio could not be computed.



Weekly Geosynchronous Satellite Environment Summary Week Beginning 02 November 2009

# GOES-11 designated Primary Electron Satellite and GOES-10 Secondary: December 1, 2008 the GOES-12 Electron sensor began experiencing periods of noise and sensor is unreliable.

Protons plot contains the five-minute averaged integral proton flux (protons/cm<sup>2</sup>-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm<sup>2</sup>-sec -sr) with energies greater than 2 MeV at GOES-11.

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-11. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWPC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





#### Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/ $m^2$ ) as measured by GOES 10 (W060) and GOES 11 (W135) in two wavelength bands, .05 - .4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm<sup>2</sup> –sec-sr) as measured by GOES-11 for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm<sup>2</sup>-sec-sr) at greater than 10 MeV.

