Space Weather Highlights 16 November – 22 November 2009

SWO PRF 1786 24 November 2009

Solar activity was very low throughout the summary period. There was only one flare, a B1 at 1259 UTC on 17 November from a spotless plage region that later emerged as Region 1033 (N21, L=191, class/area Bxo/010 on 18 November). The week began with one spotted group, Region 1031 (N30, L=257, class/area Bxo/010 on 15 November) which decayed to plage on 17 November. Two new sunspot groups emerged on 18 November: Region 1032 (N17, L=218, class/area Bxo/010 on 18 November) and Region 1033. Region 1032 decayed back to plage on 21 November and Region 1033 had decayed to plage by the end of the day on 22 November. A coronal mass ejection was observed to enter the LASCO C2 field of view off the East limb at 0755 UTC on 21 November. The associated disk signature was observed along a filament channel in the southeast portion of the solar disk.

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit was at normal background levels during the period.

The geomagnetic field was predominantly quiet during the summary period. There was a small increase to quiet to unsettled levels on 21 November. Solar wind observations from the ACE spacecraft showed a sector boundary change from towards (negative) to away (positive) early on 18 November, followed by a change back to towards (negative) on 19 November. A co-rotating interaction region was observed around 1600 UTC on 19 November (peak density ~16 p/cc) which was followed by a gradual increase in solar wind velocity from 340 km/s up to 575 km/s between 0200 UTC on 20 November through 1900 UTC on 21 November. Solar wind velocity slowly declined through the remainder of 21-22 November.

Space Weather Outlook

25 November 2009 – 21 December 2009

Solar activity is expected to be predominantly very low with just a slight chance for brief periods of low.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at normal background levels through the period.

The geomagnetic field is expected to be quiet for 25 November. An increase to quiet to unsettled levels is expected on 26 November due to recurrence. Activity levels are expected to be predominantly quiet for the remainder of the outlook period from 26 November through 21 December.



	Radio	Sun	Sunspot	X-ray				Flares				
	Flux	spot	Area	Background	X	-ray F	lux		0	ptical		
Date	10.7 cm	No.	(10 ⁻⁶ hemi.)	С	М	Х	S	1	2	3	4
16 Novembe	er 76	12	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
17 Novembe	er 77	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
18 Novembe	er 76	29	20	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
19 Novembe	er 77	30	20	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
20 Novembe	er 76	31	20	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
21 Novembe	er 76	14	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
22 Novembe	er 76	13	10	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0

Daily Solar Data

Daily Particle Data

	Prot	on Fluence	Electron Fluence	
	(proton	ns/cm ² -day-sr)	(electrons/cm ² -day-sr)	
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV
16 November	3.5e+05	2.0e+04	4.3e+03	4.8e+04
17 November	3.5e+05	2.0e+04	4.4e+03	5.4e+04
18 November	4.5e+05	2.0e+04	4.8e+03	4.7e+04
19 November	6.5e+05	1.9e+04	4.4e+03	6.8e+04
20 November	4.4e+05	1.9e+04	4.3e+03	5.3e+04
21 November	4.7e+05	1.9e+04	4.1e+03	4.0e+04
22 November	5.6e+05	2.0e+04	4.3e+03	4.9e+04

Daily Geomagnetic Data

	Middle Latitude]	High Latitude	I	Estimated
	Fredericksburg			College]	Planetary
Date	Α	K-indices	Α	K-indices	Α	K-indices
16 November	0	0-0-0-0-0-0-0-0	0	0-0-0-0-0-0-0-0	1	0-0-0-0-1-1-0-0
17 November	0	0-0-0-0-0-0-0-0	0	0-0-0-0-0-0-0-0	1	0-0-0-0-1-0-0
18 November	2	0-0-1-1-1-0-0-1	2	0-0-2-2-0-0-0-0	2	0-0-1-1-0-0-0-1
19 November	0	0-0-0-0-0-0-1	0	0-0-0-0-0-0-0-0	2	0-1-0-0-0-0-2
20 November	2	1-0-0-0-1-1-1-1	0	0-0-0-0-0-1-0	1	1-0-0-0-0-0-1
21 November	6	3-2-1-1-1-2-1	6	1-1-3-3-2-0-1-1	8	3-1-2-2-1-0-3-2
22 November	2	2-0-1-0-1-1-0-0	3	2-1-2-0-1-2-0-0	4	2-0-1-0-1-1-0-0

	Alerts and Warnings Issued	
Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
No Alerts Issued		





	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
25 Nov	75	5	2	09 Dec	72	5	2
26	75	7	2	10	72	5	2
27	73	5	2	11	75	5	2
28	71	5	2	12	75	5	2
29	70	5	2	13	75	5	2
30	70	5	2	14	75	5	2
01 Dec	70	5	2	15	75	5	2
02	70	5	2	16	75	5	2
03	70	5	2	17	75	5	2
04	70	5	2	18	75	5	2
05	70	5	2	19	75	5	2
06	70	5	2	20	75	5	2
07	70	5	2	21	75	5	2
08	72	5	2				



	Energetic Events														
	Ti	ime		X	-ray	Opti	cal Information	ı	Peak	Sweep Freq					
Date			1⁄2		Integ	Imp/	Location	Rgn	Radio Flux	Intensity					
	Begin	Max	Max	Class	Flux	Brtns	Lat CMD	#	245 2695	II IV					
No Events	Obser	ved													

			Flare List			
					Optical	
	Time		X-ray	Imp /	Location	Rgn
Begin	Max	End	Class.	Brtns	Lat CMD	
No Fla	res Obser	rved				
1256	1259	1302	B1.5			
No Fla	res Obser	rved				
No Fla	res Obser	rved				
No Fla	res Obser	rved				
No Fla	res Obser	rved				
No Fla	res Obser	rved				
	Begin No Fla 1256 No Fla No Fla No Fla No Fla	TimeBeginMaxNo Flares Obser125612561257No Flares ObserNo Flares Obser	TimeBeginMaxEndNo Flares Observed125612591302125612591302No Flares ObservedNo Flares Observed	Flare ListTimeX-rayBeginMaxEndClass.No Flares ObservedClass.125612591302B1.5No Flares ObservedB1.5B1.5No Flares ObservedClass.No Flares ObservedClass.	Flare ListTimeX-rayImp/BeginMaxEndClass.BrtnsNo Flares Observed125612591302B1.5Imp/No Flares ObservedNo Flares ObservedImp/Imp/Imp/No Flares ObservedImp/Imp/Imp/Imp/No Flares ObservedImp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/Imp/I	Flare ListOpticalTimeX-rayImp /LocationBeginMaxEndClass.BrtnsLat CMDNo Flares ObservedB1.512591302B1.5No Flares ObservedB1.5No Flares ObservedNo Flares Observed

	Region Summary															
-	Location	n		Sunspot	Character	ristics						Flaı	es			
		Helio	Area	Extent	Spot	Spot	Mag		X-ra	ıy	. —		Optio	cal		
Date (°	Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	С	Μ	Х	S	1	2	3	4	
Region 1031																
15 Nov	N30E20	258	10	1	AXX	1	А									
16 Nov	N30E08	257	10	3	BXO	2	В									
17 Nov	N30W05	5 257														
18 Nov	N30W18	3 257														
19 Nov	N30W31	257														
20 Nov	N30W44	4 257														
21 Nov	N30W57	7 257														
22 Nov	N30W7() 257														
								0	0	0	0	0	0	0		0
Still on	Disk.															

Absolute heliographic longitude: 257



			Re	egion S	lummar	y - con	tinued								
	Locatio	<u>n</u>		Sunspot	Character	ristics					I	Flare	es		
		Helio	Area	Extent	Spot	Spot	Mag		X-ra	у		(Optic	al	
Date ((° Lat ° CMD)	Lon	(10^{-6} hemi)	(helio)	Class	Count	Class	С	Μ	Х	S	1	2	3	4
	Re	gion 10)32												
18 Nov	v N17E20	218	10	2	BXO	4	В								
19 Nov	v N15E07	218	10	2	BXO	3	В								
20 No	v N16W03	3 218	10	2	BXO	2	В								
21 No	v N16W10	0 209													
22 No	v N16W2	3 209													
								0	0	0	0	0	0 ()	0
								0	Ŭ	0	Ŭ	0	<u> </u>	0	0
Still or	n Diek														
Still or	1 Disk. 1te beliogra	nhic lo	ngitude: 219	2											
Still or Absolu	n Disk. 1te heliogra	phic lo	ngitude: 218	8											
Still oı Absolı	n Disk. 1te heliogra <i>Re</i>	phic lo	ngitude: 218 9 <i>33</i>	8											
Still or Absolu 18 Nov	n Disk. ate heliogra <i>Re</i> v N21E47	phic lo gion 10 191	ngitude: 218 9 <i>33</i> 10	8	вхо	5	В								
Still or Absolu 18 Nov 19 Nov	n Disk. ate heliogra <i>Re</i> v N21E47 v N19E34	phic lo gion 10 191 191	ngitude: 218 9 <i>33</i> 10 10	8 3 5	BXO BXO	5 7	B B								
Still or Absolu 18 Nov 19 Nov 20 Nov	n Disk. ate heliogra <i>Re</i> v N21E47 v N19E34 v N21E23	phic los gion 10 191 191 189	ngitude: 218 9 <i>33</i> 10 10 10	8 3 5 1	BXO BXO BXO	5 7 9	B B B								
Still or Absolu 18 Nov 19 Nov 20 Nov 21 Nov	n Disk. ate heliogra <i>Re</i> v N21E47 v N19E34 v N21E23 v N18E07	phic los gion 10 191 191 189 192	ngitude: 218 933 10 10 10 10	8 3 5 1	BXO BXO BXO BXO	5 7 9	B B B								
Still or Absolu 18 Nov 19 Nov 20 Nov 21 Nov	n Disk. ate heliogra <i>Re</i> v N21E47 v N19E34 v N21E23 v N18E07	phic los gion 10 191 191 189 192	ngitude: 218 933 10 10 10 10 10	8 3 5 1 4	BXO BXO BXO BXO	5 7 9 4	B B B B								
Still or Absolu 18 Nov 19 Nov 20 Nov 21 Nov 22 Nov	n Disk. ate heliogra <i>Re</i> v N21E47 v N19E34 v N21E23 v N18E07 v N18W03	phic los gion 10 191 191 189 192 5 191	ngitude: 218 933 10 10 10 10 10 10	3 5 1 4 6	BXO BXO BXO BXO BXO	5 7 9 4 3	B B B B B								
Still or Absolu 18 Nov 19 Nov 20 Nov 21 Nov 22 Nov	n Disk. ate heliogra v N21E47 v N19E34 v N21E23 v N18E07 v N18W03	phic los gion 10 191 191 189 192 5 191	ngitude: 218 9 <i>33</i> 10 10 10 10 10 10	3 5 1 4 6	BXO BXO BXO BXO BXO	5 7 9 4 3	B B B B B	0	0	0	0	0	0 (0	0
Still or Absolu 18 Nov 19 Nov 20 Nov 21 Nov 22 Nov Still or	n Disk. ate heliogra v N21E47 v N19E34 v N21E23 v N18E07 v N18W0: n Disk.	phic los gion 10 191 191 189 192 5 191	ngitude: 218 9 <i>33</i> 10 10 10 10 10	3 5 1 4 6	BXO BXO BXO BXO BXO	5 7 9 4 3	B B B B	0	0	0	0	0	0 (0	0



		0		observeu l	nominiy	mean values	T 1	0	
	~ .	Sunsp	ot Numbe	rs		Radio	Flux	Geoma	gnetic
	Observed	values	<u>Ratio</u>	Smooth	values	*Penticton	Smooth	Planetary	Smooth
Month	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value
November	2.5	1.7	0.68	9.4	5.7	69.6	71.4	7	7.8
December	16.2	10.1	0.62	8.1	4.9	78.5	70.8	6	7.8
				2	008				
January	5.1	3.3	0.65	6.9	4.2	74.3	70.3	8	7.8
February	3.8	2.1	0.55	5.9	3.6	71.1	69.9	11	7.6
March	15.9	9.3	0.58	5.3	3.3	72.9	69.8	11	7.5
April	4.9	2.9	0.59	5.3	3.4	70.2	69.8	9	7.3
May	5.7	3.2	0.56	5.7	3.5	68.4	69.8	6	7.2
June	4.2	3.4	0.81	5.2	3.3	65.9	69.4	7	7.0
July	1.0	0.8	0.80	4.5	2.8	65.7	68.8	5	6.8
August	0.0	0.5	**	4.4	2.7	66.3	68.6	5	6.3
September	r 1.5	1.1	0.73	3.7	2.3	67.1	68.4	6	5.8
1									
October	5.2	2.9	0.56	2.9	1.9	68.3	68.2	7	5.4
November	6.8	4.1	0.60	2.7	1.8	68.6	68.3	4	5.1
December	1.3	0.8	0.62	2.7	1.7	69.2	68.5	4	4.9
				2	009				
January	2.8	1.3	0.46	3.0	1.8	69.8	68.7	4	4.7
February	2.5	1.4	0.56	3.1	1.9	70.0	68.8	5	4.7
March	0.7	0.7	1.00	3.4	2.0	69.2	69.0	5	4.6
April	1.2	1.2	1.00			69.7		4	
May	3.9	2.9	0.74			70.5		4	
June	6.6	2.6	0.39			68.6		4	
								-	
July	5.0	3.5	0.70			68.2		4	
August	0.3	0.0	0.00			67.4		5	
September	r 6.6	4.2	0.64			70.5		4	
~								•	
October	7.0	4.6	0.66			72.4		3	

Recent Solar Indices (preliminary) Of the observed monthly mean values

NOTE: Values are final except for the most recent 6 months which are considered preliminary. Cycle 23 started in May 1996 with an RI=8.0. Cycle 23 maximum was April 2000 with an RI=120.8.

** SWPC sunspot number was zero, so a ratio could not be computed.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 16 November 2009

GOES-11 designated Primary Electron Satellite and GOES-10 Secondary: December 1, 2008 the GOES-12 Electron sensor began experiencing periods of noise and sensor is unreliable.

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.

Electrons plot contains the five-minute averaged integral electron flux (electrons/cm²-sec -sr) with energies greater than 2 MeV at GOES-11.

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-11. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWPC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (Watts/ m^2) as measured by GOES 10 (W060) in two wavelength bands, 0.05 - 0.4 and 0.1 - 0.8 nm. The letters A, B, C, M and X refer to x-ray event levels for the 0.1 - 0.8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm²–sec-sr) as measured by GOES-11 for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

