Space Weather Highlights 23 November – 29 November 2009

Solar activity was very low. No flares occurred during the period. The solar disk was spotless during the period.

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit was at normal background levels during the period.

The geomagnetic field was initially quiet. An increase to quiet to unsettled with some isolated high latitude active periods (as well as a single minor storm period at high latitudes) occurred on 25-26 November. Activity returned to quiet levels for the remainder of the period from 27-29 November. Solar wind observations from the ACE spacecraft showed the onset of a co-rotating interaction region mid-day on 24 November, which was followed by the onset of a high speed stream. The high speed stream continued through 25 November with peak velocities around 470 km/s, and slowly declined on 26 November. The high speed stream was associated with the extension of the southern polar coronal hole as observed in solar EUV imagery.

Space Weather Outlook 02 December 2009 – 28 December 2009

Solar activity is expected to be predominantly very low with just a slight chance for brief periods of low.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at normal background levels through the period.

The geomagnetic field is expected to be quiet for 2-4 December. An increase to quiet to unsettled levels is expected on 5-6 December due to a recurrent high speed stream. Activity levels are expected to be predominantly quiet for 7-17 December, followed by another increase to quiet to unsettled on 18 December, again due to a recurrent high speed stream. Quiet levels are expected to predominate for the remainder of the outlook period.



	Radio	Sun	Sunspot	X-ray	_			Flares				
	Flux	spot	Area	Background	X	-ray F	lux		0	ptical		
Date	10.7 cm	No.	(10 ⁻⁶ hemi.))	С	Μ	Х	S	1	2	3	4
23 Novembe	r 76	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
24 Novembe	r 75	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
25 Novembe	r 74	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
26 Novembe	r 75	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
27 Novembe	r 74	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
28 Novembe	er 73	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0
29 Novembe	r 72	0	0	<a1.0< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></a1.0<>	0	0	0	0	0	0	0	0

Daily Solar Data

Daily Particle Data


		on Fluence s/cm ² -day-sr)		Electron Fluence (electrons/cm ² -day-sr)
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV
23 November	6.7e+05	1.9e+04	4.3e+03	5.3e+04
24 November	9.9e+05	2.1e+04	4.4e+03	5.1e+04
25 November	4.9e+05	1.9e+04	3.9e+03	2.1e+05
26 November	8.2e+05	2.0e+04	4.4e+03	1.1e+06
27 November	2.6e+05	2.0e+04	4.2e+03	4.4e+05
28 November	3.2e+05	2.0e+04	4.2e+03	6.3e+05
29 November	2.9e+05	2.0e+04	4.4e+03	7.6e+05

Daily Geomagnetic Data

	N	Middle Latitude		High Latitude		Estimated
]	Fredericksburg		College]	Planetary
Date	Α	K-indices	Α	K-indices	Α	K-indices
23 November	0	0-0-0-0-1-0-0	1	0-0-0-1-1-0-0-0	0	0-0-0-0-0-0-0-0
24 November	3	0-0-0-2-1-2-2	5	0-0-1-1-3-1-2-2	6	0-0-0-1-2-2-2-3
25 November	4	2-2-2-1-1-0-0-1	10	2-1-4-4-3-0-0-0	5	3-2-2-1-1-0-0-1
26 November	3	1-0-1-2-1-1-1-1	14	0-0-4-5-3-3-2-1	6	1-0-1-2-2-2-2-2
27 November	0	0-0-1-0-0-0-0-0	1	0-0-1-1-0-0-0-0	2	0-1-1-1-0-0-0-0
28 November	2	0-0-2-1-0-1-1-0	1	0-0-1-1-0-0-0-0	3	0-0-1-1-1-1-1
29 November	1	0-0-1-0-0-1-0-0	0	0-0-0-0-0-0-0-0	1	0-0-0-0-1-0-1

	Alerts and Warnings Issued	!
Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
No Alerts Issued		





Data	Radio Flux	•	U	Data	Radio Flux		0
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm		Kp Index
02 Dec	70	5	2	16 Dec	75	5	2
03	70	5	2	17	75	5	2
04	70	5	2	18	75	8	3
05	70	8	3	19	75	5	2
06	70	8	3	20	75	5	2
07	70	5	2	21	75	5	2
08	72	5	2	22	75	5	2
09	72	5	2	23	75	5	2
10	72	5	2	24	74	5	2
11	75	5	2	25	73	5	2
12	75	5	2	26	72	5	2
13	75	5	2	27	71	5	2
14	75	5	2	28	70	5	2
15	75	5	2				



					Energe	etic Events			
	Т	ime		X-ray	OI	otical Information	n	Peak	Sweep Freq
Date			1⁄2	Int	eg Imp/	Location	Rgn	Radio Flux	Intensity
	Begin	Max	Max	Class Flu	ux Brtns	Lat CMD	#	245 2695	II IV
No Event.	s Obser	ved							

		Flare List			
				Optical	
	Time	X-ray	Imp /	Location	Rgn
Date	Begin Max End	Class.	Brtns	Lat CMD	
23 November	No Flares Observed				
24 November	No Flares Observed				
25 November	No Flares Observed				
26 November	No Flares Observed				
27 November	No Flares Observed				
28 November	No Flares Observed				
29 November	No Flares Observed				

				Re	gion Su	ımmar	y									
	Locatio	n		Sunspot	Characte	ristics						Fla	es			
		Helio	Area	Extent	Spot	Spot	Mag		X-ra	y	. –		Optio	cal		
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	С	М	Х	S	1	2	3	4	
	Re	gion 10)31													
15 Nov	v N30E20	258	10	1	AXX	1	А									
16 Nov	v N30E08	257	10	3	BXO	2	В									
17 Nov	v N30W05	5 257														
18 Nov	v N30W18	3 257														
19 Nov	v N30W31	1 257														
20 Nov	v N30W44	4 257														
21 Nov	v N30W57	7 257														
22 Nov	v N30W70) 257														
23 Nov	v N30W83	3 257														
								0	0	0	0	0	0	0		0
Crosse	d West Lin	nb.														

Absolute heliographic longitude: 257



	- ·		Re		Summa		tinued									
-	Location	n Helio	A #20		t Characte		Mag	. <u> </u>	V			Fla		iaal		
Date (°	Lat ° CMD)	Lon	Area (10 ⁻⁶ hemi)	Extent (helio)	Spot Class	Spot Count	Mag Class	\overline{C}	X-ra M	x	- s	1	Opt 2		4	
Dute	,	gion 10		(neno)	Chubb	count	Chubb	U		11	5	1	-	5		
18 Nov	N17E20	0	10	2	BXO	4	В									
	N15E07		10	2	BXO	3	В									
	N16W03		10	2	BXO	2	В									
21 Nov	N16W10) 209														
22 Nov	N16W23	3 209														
23 Nov	N16W36	5 209														
24 Nov	N16W49	9 209														
25 Nov	N16W62	2 209														
26 Nov	N16W75	5 209														
27 Nov	N16W88	3 209														
								0	0	0	0	0	0	0		0
Crossed	West Lin	ıb.														
Absolut	e heliogra	phic lo	ongitude: 21	8												
	Re	gion 10	033													
18 Nov	N21E47	191	10	3	BXO	5	В									
19 Nov	N19E34	191	10	5	BXO	7	В									
20 Nov	N21E23	189	10	1	BXO	9	В									
21 Nov	N18E07	192	10	4	BXO	4	В									
22 Nov	N18W05	5 191	10	6	BXO	3	В									
23 Nov	N18W18	3 191														
24 Nov	N18W31	191														
25 Nov	N18W44	191														
26 Nov	N18W57	7 191														
	N18W7(
28 Nov	N18W83	3 191														
								0	0	0	0	0	0	0		0
	West Lin															
A1 1 /	a halianna	mbia la	ngitude: 19	1												




					nonthly	mean values						
		Sunsp	ot Numbe	rs		Radio	Flux	Geomagnetic				
	Observed	values	<u>Ratio</u>	Smooth	values	*Penticton	Smooth	Planetary	Smooth			
Month	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value			
November	r 2.5	1.7	0.68	9.4	5.7	69.6	71.4	7	7.8			
December	16.2	10.1	0.62	8.1	4.9	78.5	70.8	6	7.8			
				2	008							
January	5.1	3.3	0.65	6.9	4.2	74.3	70.3	8	7.8			
February	3.8	2.1	0.55	5.9	3.6	71.1	69.9	11	7.6			
March	15.9	9.3	0.58	5.3	3.3	72.9	69.8	11	7.5			
April	4.9	2.9	0.59	5.3	3.4	70.2	69.8	9	7.3			
May	5.7	3.2	0.56	5.7	3.5	68.4	69.8	6	7.2			
June	4.2	3.4	0.81	5.2	3.3	65.9	69.4	7	7.0			
July	1.0	0.8	0.80	4.5	2.8	65.7	68.8	5	6.8			
August	0.0	0.5	**	4.4	2.7	66.3	68.6	5	6.3			
September	r 1.5	1.1	0.73	3.7	2.3	67.1	68.4	6	5.8			
October	5.2	2.9	0.56	2.9	1.9	68.3	68.2	7	5.4			
November	r 6.8	4.1	0.60	2.7	1.8	68.6	68.3	4	5.1			
December	1.3	0.8	0.62	2.7	1.7	69.2	68.5	4	4.9			
				2	009							
January	2.8	1.3	0.46	3.0	1.8	69.8	68.7	4	4.7			
February	2.5	1.4	0.56	3.1	1.9	70.0	68.8	5	4.7			
March	0.7	0.7	1.00	3.4	2.0	69.2	69.0	5	4.6			
April	1.2	1.2	1.00			69.7		4				
May	3.9	2.9	0.74			70.5		4				
June	6.6	2.6	0.39			68.6		4				
July	5.0	3.5	0.70			68.2		4				
August	0.3	0.0	0.00			67.4		5				
September		4.2	0.64			70.5		4				
October	7.0	4.6	0.66			72.4		3				

Recent Solar Indices (preliminary) Of the observed monthly mean values

NOTE: Values are final except for the most recent 6 months which are considered preliminary. Cycle 23 started in May 1996 with an RI=8.0. Cycle 23 maximum was April 2000 with an RI=120.8.

** SWPC sunspot number was zero, so a ratio could not be computed.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 23 November 2009

GOES-11 designated Primary Electron Satellite and GOES-10 Secondary: December 1, 2008 the GOES-12 Electron sensor began experiencing periods of noise and sensor is unreliable.

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm²-sec -sr) with energies greater than 2 MeV at GOES-11.

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-11. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWPC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (Watts/m²) as measured by GOES 10 (W060) in two wavelength bands, 0.05 - 0.4 and 0.1 - 0.8 nm. The letters A, B, C, M and X refer to x-ray event levels for the 0.1 - 0.8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-11 for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

