Solar activity was at very low levels from 10-13 January. Rgn 1140 (N34, L=189, class/area Hsx/210 on 04 January) was responsible for the majority of the activity during this period, producing numerous B-class events. Activity increased to low levels during 14 - 15 January. Three C1 flares were observed from behind the NE limb during this 2-day period, all associated with returning Region 1138 (N13, L=324). Also associated with this recent activity were 4 CMEs, all of which were directed well away from Earth. Late in the period, old Region 1138 finally rotated onto the disk and was numbered 1147 (N24, L=339, class/area Cso/150 on 16 January).

No proton events were observed at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit was at high levels from 10 - 13 and 16 January and at moderate levels 14 - 16 January.

Geomagnetic field activity ranged from quiet to active levels during the period. Quiet to active conditions prevailed 10 - 11 January due to a recurrent coronal hole high speed stream (CH HSS). Activity decreased to mostly quiet levels from 12 - 13 January. From 14 - 15 January, a second CH HSS became geoeffective and increased activity to quiet to active levels. During the summary period, solar wind velocities ranged from a low of about 450 km/s late on 13 January to a high of near 600 km/s late on 09 January. The Bz component of the interplanetary magnetic field generally varied between +/- 5 nT through the period.

Space Weather Outlook 19 January – 14 February 2011

Solar activity is expected to be at very low levels with a chance for low levels during the forecast period.

No proton events are expected at geosynchronous orbit.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be below threshold at the beginning of the forecast period. High levels are expected on 04-09 February. Flux is then expected to decrease below threshold through 14 February.

Geomagnetic field activity is expected to be at quiet to unsettled levels during 19-20 January due to a recurrent CH HSS. Recurrent coronal hole high speed streams are also expected on 02-04 February and on 09-10 February, bringing predominantly unsettled conditions during those time frames. Mostly quiet levels are expected otherwise.



## Daily Solar Data

	Radio	Sun	Sunspot	X-ray	_			Flares				
	Flux	spot	Area	Background	X	ray Fl	lux		O	ptical		
Date	10.7 cm	No.	(10 <sup>-6</sup> hemi.)	)	С	M	X	S	1	2	3	4
10 January	83	35	140	A7.2	0	0	0	0	0	0	0	0
11 January	83	26	130	A6.5	0	0	0	0	0	0	0	0
12 January	80	23	80	A6.8	0	0	0	0	0	0	0	0
13 January	80	14	10	A7.3	0	0	0	0	0	0	0	0
14 January	79	11	0	A9.3	2	0	0	0	0	0	0	0
15 January	80	11	120	B1.0	1	0	0	0	0	0	0	0
16 January	80	15	120	A7.3	0	0	0	0	0	0	0	0

## Daily Particle Data

		ton Fluence	)	Electron Fluence (electrons/cm²-day-sr)
Date	>1 MeV	>10 MeV	>100 MeV	>.6 MeV >2MeV >4 MeV
10 January	4.8e + 05	1.4e + 04	3.5e+03	8.0e + 07
11 January	3.8e + 05	1.4e + 04	3.4e+03	1.1e+08
12 January	9.1e + 05	1.4e + 04	3.3e+03	1.4e + 08
13 January	7.9e + 05	1.4e + 04	3.4e+03	8.2e+07
14 January	3.8e + 05	1.3e+04	3.1e+03	1.9e+07
15 January	2.7e + 05	1.4e + 04	3.3e+03	3.2e+07
16 January	6.8e+05	1.4e+04	3.5e+03	4.6e+07

Daily Geomagnetic Data


N	Iiddle Latitude	]	High Latitude		<u>Estimated</u>
<u> </u>			College		Planetary
A	K-indices	Α	A K-indices		K-indices
4	3-1-1-1-1-0-1-0	7	2-1-3-3-3-0-1-0	5	3-1-1-1-0-1-1
5	0-2-0-2-2-1-2	7	0-1-0-0-3-4-2-1	5	1-2-0-0-2-2-2-3
5	1-2-1-2-1	5	1-1-1-3-3-0-1-0	6	2-2-1-2-1-1-1
4	2-1-0-0-2-1-2-2	7	0-1-0-1-3-3-2-3	6	2-1-0-0-1-2-2-3
6	2-1-2-2-2-1-1	13	1-1-4-2-4-4-2-1	7	1-2-2-1-2-3-1-2
2	0-2-1-1-2-1-1-0	6	0-1-1-2-4-2-0-0	5	1-2-1-1-2-1-1
3	1-1-0-0-2-2-1-0	2	1-0-0-0-1-1-1-1	3	1-0-0-0-2-1-1-1
	4 5 5 4 6 2	4 3-1-1-1-0-1-0 5 0-2-0-2-2-2-1-2 5 1-2-1-2-1-1-2-1 4 2-1-0-0-2-1-2-2 6 2-1-2-2-2-1-1 2 0-2-1-1-2-1-1-0	Fredericksburg A K-indices A 4 3-1-1-1-1-0-1-0 7 5 0-2-0-2-2-2-1-2 7 5 1-2-1-2-1-1-2-1 5 4 2-1-0-0-2-1-2-2 7 6 2-1-2-2-2-1-1 13 2 0-2-1-1-2-1-0 6	Fredericksburg A K-indices  A K-indices  A K-indices  A K-indices  4 3-1-1-1-0-1-0 7 2-1-3-3-3-0-1-0 5 0-2-0-2-2-2-1-2 7 0-1-0-0-3-4-2-1 5 1-2-1-2-1-1-2-1 5 1-1-1-3-3-0-1-0 4 2-1-0-0-2-1-2-2 7 0-1-0-1-3-3-2-3 6 2-1-2-2-2-1-1 13 1-1-4-2-4-4-2-1 2 0-2-1-1-2-1-0 6 0-1-1-2-4-2-0-0	Fredericksburg         College           A         K-indices         A           K-indices         A           K-indices         A           K-indices         A           5         0-2-0-2-2-2-1-2         7         0-1-0-0-3-4-2-1         5           5         1-2-1-2-1-1-2-1         5         1-1-1-3-3-0-1-0         6           4         2-1-0-0-2-1-2-2         7         0-1-0-1-3-3-2-3         6           6         2-1-2-2-2-1-1         13         1-1-4-2-4-4-2-1         7           2         0-2-1-1-2-1-1-0         6         0-1-1-2-4-2-0-0         5




## Alerts and Warnings Issued

Date & Time of	Issue Type of Alert or Warning	Date & Time of Event UTC
10 Jan 1221	CONTINUED ALERT: Electron 2MeV Integral Flux >= 1000pfu	<sup>3</sup> 08 Jan 1505
11 Jan 0853	CONTINUED ALERT: Electron 2MeV Integral Flux >= 1000pfu	6 08 Jan 1505
12 Jan 1033	CONTINUED ALERT: Electron 2MeV Integral Flux >= 1000pfu	<sup>3</sup> 08 Jan 1505
13 Jan 0941	CONTINUED ALERT: Electron 2MeV Integral Flux >= 1000pfu	08 Jan 1505
16 Jan 1341	ALERT: Electron 2MeV Integral Flux >= 1000pfu	16 Jan 1340









_	Radio Flux	•	_	_	Radio Flux	•	_
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
19 Jan	82	5	2	02 Feb	85	7	2
20	84	5	2	03	84	7	2
21	84	5	2	04	84	7	2
22	84	5	2	05	83	5	2
23	82	5	2	06	83	5	2
24	82	5	2	07	82	5	2
25	82	5	2	08	80	5	2
26	82	5	2	09	80	5	2
27	88	5	2	10	80	8	3
28	88	5	2	11	80	5	2
29	88	5	2	12	80	5	2
30	88	5	2	13	80	5	2
31	87	5	2	14	80	5	2
01 Feb	85	5	2				



Energetic Events

	Time	;	X-ray	Opt	ical Information	1	Peak	Sweep Freq
Date		1/2	Integ	Imp/	Location	Rgn	Radio Flux	Intensity
	Begin M	ax Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV

No Events Observed

Flare List

				Optical						
		Time		X-ray	Imp/	Location	Rgn			
Date	Begin	Max	End	Class.	Brtns	Lat CMD				
10 January	0845	0849	0852	B2.0			1145			
	0858	0904	0906	B2.5						
	1000	1028	1039	B4.6			1140			
	1343	1347	1349	B2.3			1140			
	1916	1921	1927	B1.7			1140			
	2042	2106	2112	B2.5			1140			
11 January	0606	0615	0622	B1.4			1146			
	1115	1121	1128	B1.4						
	1743	1750	1809	B3.1			1142			
	1846	1852	1901	B1.6			1142			
12 January	0039	0042	0044	B1.1			1145			
	0323	0327	0331	B1.0			1145			
	2119	2141	2205	B1.6			1140			
	2340	2355	0014	B2.2			1140			
13 January	0748	0752	0757	B1.4						
-	0909	0919	0926	B3.7			1140			
	0949	1142	1213	B2.8						
	1253	1307	1315	B2.2			1140			
	1436	1502	1514	B1.9			1140			
	2347	0006	0022	B2.3						
14 January	0128	0330	0438	C1.6						
•	1252	1307	1325	C1.0						
	1541	1551	1557	B2.2						
	1703	1709	1717	B2.2						
	1818	1824	1827	B1.8						
	2237	2258	2318	B1.8						



Flare List - continued

						Optical	
		Time		X-ray	Imp/	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	
15 January	0004	0014	0031	B2.7			
	0557	0601	0605	B2.3			
	0612	0616	0618	B2.6			
	0637	0641	0644	B4.1			
	0802	0806	0809	B3.0			
	0936	0939	0942	B3.7			
	1025	1028	1032	B2.8			
	1222	1225	1229	B2.2			
	1328	1336	1339	B2.3			
	1413	1419	1423	C1.1			
	2109	2115	2120	B2.7			1147
	2232	2238	2245	B3.3			1147
	2343	2348	2352	B5.7			1147
16 January	1237	1241	1244	B1.5			1147
· ·	1541	1544	1546	B2.4			1147
	1841	1845	1851	B1.6			1147
	1039	1048	1111	B1.4			1147
			$R \rho \sigma$	ion Summary	y		

Region Summary

	Locatio	n		Sunspot	Characte	ristics		Flares
		Helio	Area	Extent	Spot	Spot	Mag	X-ray Optical
Date (	° Lat ° CMD)	Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C M X S 1 2 3 4
	Re	gion 1	140					
31 Dec	N32E67	188	180	3	HSX	1	A	
01 Jan	N32E52	190	200	4	HSX	1	Α	1
02 Jan	N32E40	189	160	3	HSX	1	Α	
03 Jan	N32E25	190	180	2	HSX	1	A	
04 Jan	N34E14	189	210	4	HSX	1	A	
05 Jan	N34E01	188	190	6	CSO	3	В	
06 Jan	N33W1	1 188	180	3	HSX	1	Α	
07 Jan	N33W24	4 187	140	2	CSO	3	В	
08 Jan	N33W36	5 186	160	2	CSO	4	В	
09 Jan	N32W49	9 185	130	2	HSX	1	A	
10 Jan	N32W6	1 184	130	4	HSX	1	Α	
11 Jan	N33W74	4 185	120	5	CSO	3	В	
12 Jan	N32W87	7 183	70	3	HSX	1	Α	
								0 0 0 1 0 0 0 0


Crossed West Limb.

Absolute heliographic longitude: 188



Region Summary - continued

	Re	gion S	ummai	y - con	tinued					
Location		Sunspot	Character	ristics				Flares		
Helio	Area	Extent	Spot	Spot	Mag	X-ray		Optic		-
Date (° Lat ° CMD) Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C M X	S	1 2 :	3 4	
Region 1	142									
01 Jan S14E33 208	40	6	DSO	5	В					
02 Jan S12E21 207	90	8	DSO	3	В					
03 Jan S13E07 208	120	10	DSI	13	В	1	1			
04 Jan S12W05 208	90	8	DSI	17	В					
05 Jan S13W18 208	40	8	CAO	14	В					
06 Jan S14W32 208	50	8	CSO	7	В					
07 Jan S14W45 208		1	AXX	2	A					
08 Jan S15W56 205	10	2	BXO	3	В					
09 Jan S17W75 212	10	6	BXO	4	В					
10 Jan S17W88 212										
						1 0 0	1	0 0 0	)	0
Crossed West Limb.										
Absolute heliographic lo	ongitude: 208	3								
Region 1	143									
07 Jan S22E16 147	20	4	DSO	4	В					
08 Jan S23E02 146	10	6	BXO	4	В					
09 Jan S23W11 146										
10 Jan S23W24 146										
11 Jan S23W37 146										
12 Jan S23W50 146										
13 Jan S23W63 146										
14 Jan S23W76 146										
15 Jan S23W89 146										
						0 0 0	0	0 0 0	)	0
Crossed West Limb.										
Absolute heliographic lo	ongitude: 146	5								
	<b>.</b>									
Region 1	144	•								
09 Jan S16W36 173		2	AXX	1	A					
10 Jan S16W49 173										
11 Jan S16W62 173										
12 Jan S16W75 173										
13 Jan S16W88 173						0 0 0	^	0 0	`	0
Constant West 1						0 0 0	Ü	0 0 (	)	0
Crossed West Limb.		_								





Region Summary - continued

	<u> </u>	gion s	ummai	y - con	unuea					
Location			Characte					Fla	res	
Helio	Area	Extent	Spot	Spot	Mag	X-ray			Optical	_
Date (° Lat ° CMD) Lon	(10 <sup>-6</sup> hemi)	(helio)	Class	Count	Class	C M X	S	1	2 3 4	
<b>.</b>	4.5									
Region 11					_					
09 Jan N15E39 97	10	4	BXO	4	В					
10 Jan N16E27 98	10	5	BXO	3	В					
11 Jan N16E14 98										
12 Jan N16E01 98										
13 Jan N16W12 98										
14 Jan N16W25 100		1	AXX	1	A					
15 Jan N16W43 100										
16 Jan N16W56 100										
						$0 \ 0 \ 0$	0	0	0 0	0
Still on Disk.										
Absolute heliographic los	ngitude: 98									
0 1	C									
Region 11	46									
10 Jan N23E46 77		1	AXX	1	A					
11 Jan N23E33 76	10	2	BXO	3	В					
12 Jan N24E19 78	10	2	BXO	2	В					
13 Jan N23E06 76	10	3	BXO	4	В					
14 Jan N23W06 76										
15 Jan N23W19 76										
16 Jan N23W32 76										
10 3411 1(23 )( 32 ) 70						0 0 0	0	0	0 0	0
Still on Disk.						0 0 0	Ü	Ü	0 0	Ü
Absolute heliographic los	ngitude: 76									
Absolute henograpine los	ingitude. 70									
Region 11	17									
15 Jan N24E80 344	120	1	HRX	1	A					
				5	A B					
16 Jan N27E65 339	120	6	DSO	3	В	0 0 0	0	0	0 0	0
Gell Dist						0 0 0	U	U	0 0	0
Still on Disk.		_								




Absolute heliographic longitude: 339

## Recent Solar Indices (preliminary) Of the observed monthly mean values

		Cunco			monunty	Radio	Eluv	Gooma	anotio
	Observed		ot Numbe		volves			Geoma	_
3.6 .1	Observed			Smooth		Penticton 10.7		Planetary	
<u>Month</u>	SEC	RI	RI/SEC	SEC	RI	10.7 cm	Value	Ap	Value
τ.	2.0	1.0	0.46		009	60.0	60. <b>7</b>	4	4.77
January	2.8	1.3	0.46	3.0	1.8	69.8	68.7	4	4.7
February	2.5	1.4	0.56	3.1	1.9	70.0	68.8	5	4.7
March	0.7	0.7	1.00	3.4	2.0	69.2	69.0	5	4.6
A	1.0	0.0	1.00	2.7	2.2	<i>c</i> 0.7	co 2	4	4.2
April	1.2	0.8	1.00	3.7	2.2	69.7	69.3	4	4.3
May	3.9	2.9	0.74	3.8	2.3	70.5	69.7	4	4.1
June	6.6	2.9	0.39	4.4	2.7	68.6	70.2	4	4.0
July	5.0	3.2	0.70	5.8	3.6	68.2	71.0	4	3.9
August	0.3	0.0	0.00	7.7	4.8	67.4	72.1	5	3.8
September		4.3	0.64	9.9	6.2	70.5	73.3	4	3.8
Беристое	0.0	1.5	0.01	7.7	0.2	70.5	73.3	•	3.0
October	7.0	4.8	0.66	11.3	7.1	72.3	74.1	3	4.1
November	r 7.7	4.1	0.55	12.4	7.6	73.6	74.5	3	4.5
December	15.7	10.8	0.68	13.6	8.3	76.8	74.9	2	4.8
				_					
	21.2	10.0	0. 62		010	04.4	<b></b>	2	<b>7</b> 0
January	21.3	13.2	0.62	14.8	9.3	81.1	75.5	3	5.0
February	31.0	18.8	0.60	16.7	10.6	84.7	76.5	5	5.1
March	24.7	15.4	0.62	19.1	12.3	83.3	77.5	5	5.3
April	11.2	8.0	0.71	21.4	14.0	75.9	78.3	10	5.5
May	19.9	8.7	0.44	23.8	15.5	73.8	79.0	8	5.7
June	17.9	13.6	0.75	25.2	16.4	72.6	79.7	7	5.8
0 0/110	2117	10.0	0.70		101.	, =	.,,,,	•	2.0
July	23.1	16.1	0.70			79.9		5	
August	28.2	19.6	0.70			79.7		8	
September	r 35.6	25.2	0.71			81.1		5	
October	35.0	23.5	0.67			81.6		6	
November		21.6	0.60			82.5		5	
December	22.0	14.5	0.66			84.3		3	

**NOTE:** Values are final except for the most recent 6 months which are considered preliminary. Cycle 23 started in May 1996 with an RI=8.0. Cycle 23 maximum was April 2000 with an RI=120.8. Solar minimum for cycle 23 was December 2008.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 10 January 2011


The proton flux plot contains the five-minute averaged integral proton flux (protons/cm<sup>2</sup>-sec -sr) as measured by GOES-13 (W75) for each of three energy thresholds: greater than 10, 50, and 100 MeV. The electron flux plot contains the five-minute averaged integral electron flux (electrons/cm<sup>2</sup>-sec -sr) with energies greater than 2 MeV at GOES-13.

The Hp plot contains the five minute averaged Hp magnetic field component in nanoteslas (nT) as measured by GOES-13. The Hp component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWPC and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots Week Beginning 10 January 2011

The x-ray plot contains five-minute averaged x-ray flux (Watts/ $m^2$ ) as measured by GOES 14 (W105) in two wavelength bands, 0.05 - 0.4 and 0.1 - 0.8 nm. The letters A, B, C, M and X refer to x-ray event levels for the 0.1 - 0.8 nm band.

The proton plot contains the five-minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-13 for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

