

RECALL FROM LAST TIME

FUNDAMENTAL THEOREM OF CALCULUS

$$\int_a^b \left(\frac{df}{dx} \right) dx = f(b) - f(a)$$

GRADIENT THEOREM

$$\int_{\vec{A}}^{\vec{B}} (\nabla T) \cdot d\vec{l} = T(\vec{B}) - T(\vec{A})$$

COROLLARIES:

- THE ABOVE INTEGRAL IS PATH INDEPENDENT.
- $\oint (\nabla T) \cdot d\vec{l} = 0$.

DIVERGENCE THEOREM (GAUSS'S THM, GREEN'S THM)

$$\int_V (\nabla \cdot \vec{v}) dV = \oint_S \vec{v} \cdot d\vec{a}$$

EXAMPLE: WHAT IS $\oint_S (\nabla \times \vec{A}) \cdot d\vec{a}$?

TODAY: STOKES' THM (CURL THEOREM) GRIFITHS § 1.3.5

$$\int_A (\nabla \times \vec{v}) \cdot d\vec{a} = \oint_C \vec{v} \cdot d\vec{l}$$


IMPORTANT: DIRECTION OF $d\vec{a} = \hat{n} da$ GIVEN BY
RIGHT HAND RULE ON CONTOUR C.

(2)

IN THE NEIGHBORHOOD OF SOME POINT \vec{r}_i ,

CONSIDER A SMALL BIT OF AREA ΔA_i ON A LARGER SURFACE Σ . WITH NORMAL VECTOR \hat{n} .


CONSTRUCT A COORDINATE SYSTEM WITHIN ΔA_i , ORIGIN AT CENTER.



$$\text{Let } \vec{v}_i = \vec{v}(\vec{r}_i).$$

ALSO, LET Σ BE SMALL ENOUGH THAT DERIVATIVES OF $\vec{v}(\vec{r})$ ARE CONSTANT. Thus $\vec{v} = \vec{v}_i + x \frac{\partial \vec{v}}{\partial x} + y \frac{\partial \vec{v}}{\partial y}$ WITHIN A_i .

WE SEEK TO COMPUTE $\int_{\partial A_i} \vec{v} \cdot d\vec{l}$, ON THE BOUNDARY OF A_i .



$$\oint_{\partial A_i} \vec{v} \cdot d\vec{l} = \int_{C_1} \vec{v}_i \cdot d\vec{l} + \int_{C_2} \left(x \frac{\partial \vec{v}}{\partial x} + y \frac{\partial \vec{v}}{\partial y} \right) \cdot d\vec{l}$$

$$C_1: \text{FOR } x = \epsilon/2, \quad d\vec{l} = \hat{y} dy, \quad \cancel{\int_{C_1} \vec{v}_i \cdot d\vec{l}}$$

$$\int_{C_1} \left(x \frac{\partial \vec{v}}{\partial x} + y \frac{\partial \vec{v}}{\partial y} \right) \cdot d\vec{l} = \int_{-\epsilon/2}^{\epsilon/2} \left(\frac{\epsilon}{2} \frac{\partial v_y}{\partial x} + y \frac{\partial v_y}{\partial y} \right) dy =$$

$$= \left[\frac{\epsilon}{2} \frac{\partial v_y}{\partial x} y + \frac{1}{2} y^2 \frac{\partial v_y}{\partial y} \right]_{-\epsilon/2}^{\epsilon/2} = \frac{\epsilon^2}{2} \frac{\partial v_y}{\partial x}.$$

(3)

$$C_2: y = \varepsilon h, \quad d\vec{r} = \hat{x} dx$$

$$\begin{aligned} \int_{C_2} \left(x \frac{\partial \vec{v}}{\partial x} + y \frac{\partial \vec{v}}{\partial y} \right) \cdot d\vec{r} &= \int_{\varepsilon h}^{-\varepsilon h} \left(x \frac{\partial v_x}{\partial x} + \frac{\varepsilon}{2} \frac{\partial v_x}{\partial y} x \right) dx \\ &= \left[\frac{1}{2} x^2 \frac{\partial v_x}{\partial x} + \frac{\varepsilon}{2} \frac{\partial v_x}{\partial y} x \right]_{\varepsilon h} \\ &= -\frac{\varepsilon}{2} \frac{\partial v_x}{\partial y}. \end{aligned}$$

$$C_3: \text{ same as } C_1, \quad \int_{C_3} = \frac{\varepsilon}{2} \frac{\partial v_y}{\partial x}.$$

$$C_4: \text{ same as } C_2, \quad \int_{C_4} = -\frac{\varepsilon}{2} \frac{\partial v_x}{\partial y}.$$


$$\therefore \oint \vec{v} \cdot d\vec{r} = \frac{\varepsilon}{2} \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right).$$

$$= (\nabla \times \vec{v}) \cdot \hat{z} A = (\nabla \times \vec{v}) \cdot \hat{n} \int_A da$$

SINCE $\nabla \times \vec{v} = \text{curl}$,

$$\oint \vec{v} \cdot d\vec{r} = \int_A (\nabla \times \vec{v}) \cdot \hat{n} da = \int_A (\nabla \times \vec{v}) \cdot d\vec{a}$$

NOW, NOTICE THAT LARGER AREAS, OF ARBITRARY SHAPE, CAN BE CONSTRUCTED BY JOINING SMALLER ONES:




The GAO & DIV THEOREMS CAN BE DEMONSTRATED SIMILARLY.

(4)

CORROLARIES OF STOKES' THEOREM

$\oint_A (\nabla \times \vec{v}) \cdot d\vec{a}$ IS INDEPENDENT OF SURFACE FOR ALL SURFACES BOUNDED BY THE SAME PATH.



$\oint (\nabla \times \vec{v}) \cdot d\vec{a} = 0$ ("SHRINKING MOUTH" ARGUMENT)

CONNECTION TO GAUSS'S THEOREM:

$$\oint (\nabla \times \vec{v}) \cdot d\vec{a} = \int_V \nabla \cdot (\nabla \times \vec{v}) dV = \int 0 dV = 0 \checkmark$$


CONNECTION TO GRADIENT THEOREM:

$$\begin{aligned} & \oint (\nabla T) \cdot d\vec{l} = 0 \quad \leftarrow \text{CORROLARY OF } \nabla \text{ THM.} \\ & = \int [\nabla \times (\nabla T)] \cdot d\vec{a} \xrightarrow{\text{STOKES}} \int \vec{0} \cdot d\vec{a} = 0 \checkmark \end{aligned}$$

EXAMPLE / EXERCISE

(5)

LET $\vec{v}(r)$ REPRESENT THE VELOCITY OF A DISK OF RADIUS R , SPINNING AT ANGULAR FREQUENCY ω .



(a) FIND $\nabla \times \vec{v}$

(b) CALCULATE $\oint \vec{v} \cdot d\vec{l}$ AROUND THE DISK'S CIRCUMFERENCE.

(c) ~~DISCUSS~~ VERIFY STOKES' THM BY CALCULATING $\int (\nabla \times \vec{v}) \cdot d\vec{a}$.

$$\vec{v} = \omega r \hat{\phi} = \omega(x\hat{y} - y\hat{x}).$$

$$(a) \nabla \times \vec{v} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -yw & xw & 0 \end{vmatrix} = 2\omega \hat{z}$$

$$(b) \oint \vec{v} \cdot d\vec{l} = \int_0^{2\pi} (r\omega \hat{\phi}) \cdot (r\hat{\phi} dr) = 2\pi r^2 \omega$$


$$(c) \int (\nabla \times \vec{v}) \cdot d\vec{a} = 2\omega \int da = 2\pi r^2 \omega \checkmark$$

(5)

EXERCISE /
EXAMPLE

$$\vec{v} = \hat{x} \cos z + \hat{y} \sin z + \hat{z}$$

VERIFY STOKES' THEOREM FOR THE ILLUSTRATED PATH.



$$C_1: \vec{v} = \hat{x} + \hat{z}, \quad d\vec{l} = \hat{y} dy$$

$$\int_{C_1} \vec{v} \cdot d\vec{l} = 0$$

$$C_2: z = \pi - y, \quad d\vec{l} = \hat{z} dz + \hat{y} dy = (\hat{z} - \hat{y}) dy$$

LET'S PUT EVERYTHING IN TERMS OF z !

$$\vec{v} \cdot d\vec{l} = (1 - \sin z) dz$$

$$\int_{C_2} \vec{v} \cdot d\vec{l} = \int_0^\pi (1 - \sin z) dz = [z + \cos z]_0^\pi = \pi - 2$$

$$C_3: y=0, \quad d\vec{l} = dz \hat{z}, \quad \vec{v} \cdot d\vec{l} = dz$$

$$\int_0^\pi dz = -\pi$$

$$\therefore \oint \vec{v} \cdot d\vec{l} = -2$$

NOW,

$$\nabla \times \vec{v} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \cos z & \sin z & 1 \end{vmatrix} = -\hat{x} \cos z - \hat{y} \sin z, \quad d\vec{n} = \hat{x} dy dz$$

$$\text{so } (\nabla \times \vec{v}) \cdot d\vec{n} = -\cos z dy dz.$$

$$\begin{aligned} \int (\nabla \times \vec{v}) \cdot d\vec{n} &= \int_{y=0}^{\pi} \left[\int_{z=0}^{\pi-y} -\cos z dz \right] dy = \int_0^{\pi} \left[-\sin z \Big|_0^{\pi-y} \right] dy \\ &= \int_0^{\pi} \sin(\pi-y) dy = -\cos(y-\pi) \Big|_0^{\pi} = -[1+1] = -2 \end{aligned}$$