

## § 24.13 — APPLYING CONTOUR INTEGRATION TO DEFINITE INTEGRALS


ONE OF THE MOST IMPORTANT APPLICATIONS IS TO  
IMPROPER INTEGRALS, E.G.

$$\int_{-\infty}^{\infty} f(x) dx$$

WE WRITE THIS AS

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx,$$

STRATEGY: CONVERT TO A CONTOUR INTEGRAL



$$\int_{-\infty}^{\infty} f(x) dx = \int_C f(z) dz - \int_{\Gamma} f(z) dz$$

↑  
 SEMICIRCLE  
 PLUS  $-R < x < R$   
 ON X-AXIS

$$= \sum_{\text{INSIDE}} R - \int_{\Gamma} f(z) dz$$


↑ HOPE THIS IS EASIER  
THAN THE ORIGINAL INTEGRAL!

EXAMPLE EVALUATE:

$$I = \int_{-\infty}^{\infty} \frac{dx}{a^2 + x^2}$$

$$\int_{-\infty}^{\infty} \frac{dx}{a^2 + x^2} = \oint_C \frac{dz}{(z+ia)(z-ia)} - \int_P \frac{dz}{a^2 + z^2}$$

WE HAVE JUST ONE SIMPLE POLE INSIDE THE CONTOUR:



$$R(i\alpha) = \lim_{z \rightarrow ia} \left\{ \frac{1}{z+ia} \right\} = \frac{1}{2ia}$$

$$I = 2\pi i \sum_{\text{INSIDE}} R = \frac{2\pi i}{2ia}$$

$$\int_{-\infty}^{\infty} \frac{dx}{a^2 + x^2} = \frac{\pi}{a}$$


(8)

CONSIDER THE F.T. INTEGRAL

$$\begin{aligned} \mathcal{F}\left(\frac{1}{1+x^2}\right) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{e^{-ikx}}{1+x^2} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_L \frac{e^{-ikz}}{1+z^2} dz \end{aligned}$$

WHERE CONTOUR L IS THE REAL AXIS.

FOR  $k > 0$ , THE CONTOUR CAN BE CLOSED WITH A SEMICIRCLE, ALONG WHICH THE CONTOUR INTEGRAL IS ZERO. SHOULD I CHOOSE THE UPPER ( $\Gamma$ ) OR LOWER ( $\Gamma'$ ) SEMICIRCLE?



EVALUATE  $\oint$ , AND HENCE FIND  $\mathcal{F}\left(\frac{1}{1+x^2}\right)$  FOR  $k > 0$ .

FIND  $\mathcal{F}\left(\frac{1}{1+x^2}\right)$  FOR  $k < 0$

COMBINE.

Complex expansion:

OUR NUMERATOR,  $e^{ikz}$ , LOOKS INNOCENT.

JUST A COMPLEX NUMBER OF UNIT MODULUS, RIGHT?

Q: WHAT'S DIFFERENT ABOUT  $e^{ikz}$ ?

A: LOOK AT  $z = r \cos \theta + ir \sin \theta$ . THEN

$$e^{-ikz} = e^{-ikr \cos \theta} e^{ikr \sin \theta}$$

↑

BLOWS UP AS  $r \rightarrow \infty$ ,  
FOR  $0 < \theta < \pi$ , UNLESS  $k < 0$ .

SO INTEGRATE ON  $\Gamma$  FOR  $k < 0$

NOW, FOR  $k > 0$ , INTEGRATE ON  $\Gamma'$  SO  $\sin \theta \leq 0$ .

THE RESULT SHOULD BE SOMETHING LIKE

$$\mathcal{I}\left[\frac{1}{x^2+1}\right] = \begin{cases} \sqrt{\frac{\pi}{2}} e^{2k}, & k < 0 \\ \sqrt{\frac{\pi}{2}} e^{-2k}, & k > 0 \end{cases}$$

$$= \sqrt{\frac{\pi}{2}} e^{-2|k|}$$

9


EXAMPLE EVALUATE

$$I = \int_{-\infty}^{\infty} \frac{dx}{(x^2 - 2x + 4)}$$

Roots:  $\lambda = \frac{2 \pm \sqrt{4-16}}{2} = 1 \pm i\sqrt{3}$

(THESE ARE SIMPLE POLES)

## Example



6. Let  $f(z) \equiv 1/(1+z^2)$ .

(a) (5 points) Where is  $f(z)$  analytic?

(b) (15 points) It is easy to show using trigonometric substitution that

$$\int_0^\infty \frac{dx}{1+x^2} = \frac{\pi}{2}.$$

Use this fact and integrate  $f(z)$  on the illustrated contour to show that

$$\begin{aligned} \int_0^\infty \frac{1+r^2}{1+r^4} dr &= \cancel{\frac{\pi}{2\sqrt{2}}} \text{ and } \frac{\pi}{\sqrt{2}} \\ \int_0^\infty \frac{1-r^2}{1+r^4} dr &= 0. \end{aligned}$$

(a)  $f(z)$  is analytic everywhere except at the roots of the denominator,  $z = \pm i$ .

$$(b) 0 = \oint f(z) dz = \int_I f(z) dz + \int_{II} f(z) dz + \int_{III} f(z) dz$$

$$0 = \int_0^\infty \frac{dx}{1+x^2} + 0 + \int_{III} f(z) dz \quad *$$

on PATH III,  $dz = e^{i\pi/4} dr$ ;  $z^2 = x^2 - y^2 + 2ixy = 2ix^2$   
 $r^2 = x^2 + y^2 = 2x^2$

$$\therefore \int_{III} f(z) dz = - \int_{r=0}^\infty \frac{e^{i\pi/4} dr}{1+i r^2}$$

cont.

scratch paper

$$\begin{aligned}\int_{\text{III}} f dz &= - \int_0^\infty \frac{\frac{1}{\sqrt{2}}(1+i)}{1+ir^2} dr \\ &= -\frac{1}{\sqrt{2}} \int_0^\infty \frac{(1+i)(1-ir^2)}{1+r^4} dr \\ &= -\frac{1}{\sqrt{2}} \int_0^\infty \frac{[(1+r^2)+i(1-r^2)]}{1+r^4} dr\end{aligned}$$

PLUGGING BACK INTO \*, THE REAL PART IS:

$$\int_0^\infty \frac{1+r^2}{1+r^4} dr = \frac{\pi}{\sqrt{2}} \quad \checkmark$$

AND IMAGINARY PART:

$$\int_0^\infty \frac{1-r^2}{1+r^4} dr = 0 \quad \checkmark$$

## JORDAN'S LEMMA

NOTE THAT  $\int_{\gamma} \rightarrow 0$  WAS VERY IMPORTANT.

WE NOW EXPAND THE SET OF SUCH SEMICIRCLES THAT INTEGRATE TO ZERO.

IF  $f(z)$  IS ANALYTIC FOR  $y > 0$  EXCEPT FOR A FINITE # POLES AND  $\max |f(z)| \rightarrow 0$  AS  $|z| \rightarrow \infty$  FOR  $y > 0$  AND  $\Im m > 0$ ,

$$I_{\Gamma} = \int_{\Gamma} e^{imz} f(z) dz \rightarrow 0 \text{ AS } R \rightarrow \infty$$


PROOF

$$I_{\Gamma} \leq \int_{\Gamma} |e^{imz}| |f(z)| |dz| \stackrel{Rd\theta}{\sim} M \int_0^{\pi} |e^{imz}| Rd\theta$$

$$\text{NOW, } |e^{imz}| = |e^{imx} e^{-my}| = e^{-my} = e^{-mR \sin \theta}$$

$$\Rightarrow I_{\Gamma} \leq MR \int_0^{\pi} e^{-mR \sin \theta} d\theta = 2MR \int_0^{\pi/2} e^{-mR \sin \theta} d\theta$$

$$\text{ON } 0 < \theta < \pi/2, \sin \theta \geq \frac{2\theta}{\pi},$$



$$\text{so } e^{-mR \sin \theta} \leq e^{-mR(2\theta/\pi)}$$

$$\therefore I_{\Gamma} \leq 2MR \int_0^{\pi/2} e^{-2mR\theta/\pi} = \frac{\pi M}{m} (1 - e^{-mR}) < \frac{\pi M}{m}$$

BUT RECALL  $M \rightarrow 0$ , SO

$$\lim_{R \rightarrow \infty} I_{\Gamma} = 0 \quad \underline{\text{QED}}$$