300 Seconds Above 160 Kilometers

Charles Kankelborg Physics Department, Montana State University

2016 OpTeC Conference







Abstract



Slitless spectrographs offer a way to measure doppler shifts across a wide field of view in a single snapshot. MOSES, a rocket borne instrument forming simultaneous images of He II Ly alpha (30.4 nm, T 50,000 K) at several spectral orders, observed explosive events in the solar atmosphere on February 8, 2006. A second launch on August 27, 2015 observed doppler shifts in a solar active region in Ne VII (46.5 nm, T 500,000 K). We are now preparing for a third flight and adding an instrument of improved design, ESIS. I will summarize what we have learned and what we expect to accomplish with ESIS.







Motivation
MOSES-I
MOSES-II
ESIS



Extreme Ultraviolet Sun





Observing From Space



• Atmosphere is opaque to EUV

- In air (1 atm), EUV propagates about 0.01 mm
- 50% transmission at 160 km altitude
- NASA Sounding Rocket Program
 - Suborbital
 - Inexpensive (1% of a satellite)
 - Risk tolerance

Right: MOSES-I launch, 2006-Feb-08







Slitless spectrograph, He II λ 304 Å. Differences between images encode the line profile.



MOSES-I Images



6



Fox, Kankelborg and Thomas (2010) Astrophys. J. 719:1132



MOSES-I Dopplergrams



Bright red and blue correspond to line of sight velocities around 50 km/s.



Courrier and Kankelborg, 2015 *Proc. SPIE* 9643, doi:10.1117/12.2194607



MOSES-I Inversions



Multiresolution inversion of spatial-spectral cube. Many explosive events in the solar transition region.



Tom Rust (to be submitted to Astrophys. J.)























A solar active region in NeVII, 46.5 nm.



Observations are rare in this temperature range (500,000 K).



EUV Snapshot Imaging Spectrometer (ESIS)



- New instrument design, flight in 2018
- Optics designed/build/test at MSU
- Detector system from NASA/MSFC (Jon Cirtain, Ken Kobayashi)



Recent Work



Afternoon Poster Session

- Roy Smart & Hans Courrier "Preliminary results of the MOSES II 2015 flight." (SPD 2016)
- Shane Atwood "A PSF equalization technique for the multi-order solar extreme ultraviolet spectrograph (MOSES)" (in preparation for JEI)

In Preparation

- Tom Rust "Transition Region Explosive Events in He II" (ApJ)
- Hans Courrier "Using local correlation tracking to obtain solar spectral information from a slitless spectrograph" (*JEI*)