Physics 423, Spring 2019

Notice: We will have an in-class exam on Monday, Jan 28:

- The exam covers Chapter 1 of Griffiths and is worth 50 points.
- There will be 3 ± 1 problems, similar to past homework.
- You may bring a 4x6 card with notes if you like. I will provide something like the formulae on the endpapers of your text.
- I will hold a review session in class Friday, Jan 25.

Homework Set # 3

- 1. Due Wednesday, Jan 23 (Read §1.5):
 - (a) In class we found that

$$\int_0^\infty f(x)\,\delta(x^2 - 2)\,dx = \frac{f(\sqrt{2})}{2\sqrt{2}}\,.$$

Verify this result by factoring the argument of the delta function, and then using the identity

$$\int_{-\infty}^{\infty} f(x)\,\delta([x-b]c)\,dx = \frac{f(b)}{|c|}.$$

- (b) Let $v(r) = (1 + \cos \theta) \hat{r} / r^2$.
 - i. Calculate $\nabla \cdot v$.
 - ii. Calculate $\oint \boldsymbol{v} \cdot d\boldsymbol{a}$ for the sphere r = R.
 - iii. If necessary, correct the divergence result with what you learned in the previous part.
- 2. Due Friday, Jan 25 (Read §1.6):
 - (a) The "vector area" of surface S is defined by $a \equiv \int_{S} da$. Show that the vector area of a closed surface is zero.¹
 - (b) Similarly, show that $\oint d\mathbf{l} = 0$.
 - (c) Use the theorems in Section 1.6 to show that $\mathbf{F} = yz\hat{\mathbf{x}} + xz\hat{\mathbf{y}} + xy\hat{\mathbf{z}}$ can be written both as the gradient of a scalar and as the curl of a vector. Then, find the scalar and vector potentials for \mathbf{F} .

¹Hint: calculate the components one at a time, for example $a_x = \hat{x} \cdot a$.